

ANALYSIS RELATED TO ALL ADMISSIBLE TYPE PARAMETERS IN THE JACOBI SETTING

ADAM NOWAK, PETER SJÖGREN, AND TOMASZ Z. SZAREK

ABSTRACT. We derive an integral representation for the Jacobi-Poisson kernel valid for all admissible type parameters α, β in the context of Jacobi expansions. This enables us to develop a technique for proving standard estimates in the Jacobi setting, which works for all possible α and β . As a consequence, we can prove that several fundamental operators in the harmonic analysis of Jacobi expansions are (vector-valued) Calderón-Zygmund operators in the sense of the associated space of homogeneous type, and hence their mapping properties follow from the general theory. The new Jacobi-Poisson kernel representation also leads to sharp estimates of this kernel. The paper generalizes methods and results existing in the literature, but valid or justified only for a restricted range of α and β .

1. INTRODUCTION

This paper is a continuation and completion of the research performed recently in [7] by the first and second authors. Given parameters $\alpha, \beta > -1$, consider the Jacobi differential operator

$$\mathcal{J}^{\alpha, \beta} = -\frac{d^2}{d\theta^2} - \frac{\alpha - \beta + (\alpha + \beta + 1) \cos \theta}{\sin \theta} \frac{d}{d\theta} + \left(\frac{\alpha + \beta + 1}{2} \right)^2$$

on the interval $[0, \pi]$ equipped with the (doubling) measure

$$d\mu_{\alpha, \beta}(\theta) = \left(\sin \frac{\theta}{2} \right)^{2\alpha+1} \left(\cos \frac{\theta}{2} \right)^{2\beta+1} d\theta.$$

This operator, acting initially on $C_c^2(0, \pi)$, has a natural self-adjoint extension in $L^2(d\mu_{\alpha, \beta})$, whose spectral decomposition is discrete and given by the classical Jacobi polynomials. Various aspects of harmonic analysis related to the Jacobi setting has been studied in the literature. This line of research goes back to the seminal work of B. Muckenhoupt and E.M. Stein [6], in which the ultraspherical case ($\alpha = \beta$) was investigated. Later several other authors contributed to the subject, see [7, Section 1] and also the end of [7, Section 2] for a detailed account and references.

The main result of [7] is restricted to $\alpha, \beta \geq -1/2$. It states that several fundamental operators in the harmonic analysis of Jacobi expansions, including Riesz transforms, imaginary powers of the Jacobi operator, the Jacobi-Poisson semigroup maximal operator and Littlewood-Paley-Stein type square functions, are (vector-valued) Calderón-Zygmund operators in the sense of the space of homogeneous type $([0, \pi], d\mu_{\alpha, \beta}, |\cdot|)$. Here $|\cdot|$ stands for the ordinary distance. Consequently, the mapping properties of these operators follow from the general theory. The proofs in [7] rely

2010 Mathematics Subject Classification: primary 42C05; secondary 42C10

Key words and phrases: Jacobi polynomial, Jacobi expansion, Jacobi operator, Jacobi-Poisson kernel, sharp estimate, Jacobi-Poisson semigroup, Riesz transform, multiplier, imaginary power, fractional integral, maximal operator, square function, Calderón-Zygmund operator.

The first author was supported in part by MNiSW Grant N N201 417839.

on an integral formula for the Jacobi-Poisson kernel, derived in [7] from a product formula for Jacobi polynomials due to Dijksma and Koornwinder [4]. Unfortunately, the latter result is not valid if either $\alpha < -1/2$ or $\beta < -1/2$, and this limitation is inherited by the abovementioned Jacobi-Poisson kernel representation. Thus the technique of proving estimates for kernels defined via the Jacobi-Poisson kernel developed in [7] is designed for the case $\alpha, \beta \geq -1/2$. The object of the present paper is to eliminate this restriction in the parameter values, which will require some new techniques.

Our method starts with the deduction of a suitable integral representation of the Jacobi-Poisson kernel, valid for all $\alpha, \beta > -1$, see Proposition 2.3. This extended Jacobi-Poisson kernel formula contains as a special case that obtained in [7, Proposition 4.1] for $\alpha, \beta \geq -1/2$ and is more involved if either α or β is less than $-1/2$. Then we establish a suitable generalization to all $\alpha, \beta > -1$ of the strategy employed in [7] to prove standard estimates (see (15)-(17) below) for kernels expressible via the Jacobi-Poisson kernel. To achieve this, some essentially new arguments are required, and the method allows a unified treatment of all parameter values $\alpha, \beta > -1$.

As an application of these techniques, we prove that the maximal operator of the Jacobi-Poisson semigroup, the Riesz-Jacobi transforms, Littlewood-Paley-Stein type square functions and multipliers of Laplace and Laplace-Stieltjes transform type are scalar-valued or vector-valued Calderón-Zygmund operators in the sense of the space $([0, \pi], d\mu_{\alpha, \beta}, |\cdot|)$; see Theorem 5.1. This extends to all $\alpha, \beta > -1$ several results for $\alpha, \beta \geq -1/2$ obtained in [7] and earlier papers. Moreover, here we treat Laplace transform type multipliers, which were considered in [7] only implicitly (see the comment closing [7, Section 2]), and also Laplace-Stieltjes transform type multipliers not studied earlier in the Jacobi context. These multiplier operators include, as special cases, the imaginary powers of the Jacobi operator and the related fractional integrals (provided that $\alpha + \beta + 1 \neq 0$; otherwise the bottom eigenvalue of the Jacobi operator is 0 and these objects are not well defined). Noteworthy, our technique is well suited to a wider variety of operators, including more general forms of g -functions and Lusin area type integrals. In a similar spirit, analogous problems concerning analysis for “low” values of type parameters were recently investigated in the Laguerre [9] and Bessel [2] settings.

The utility of the Jacobi-Poisson kernel representation derived in Proposition 2.3 is also strongly supported by the fact that it makes it possible to describe the exact behavior of the kernel; see Theorem 6.1. The sharp estimates we prove extend to all $\alpha, \beta > -1$ the bounds found recently by Nowak and Sjögren [8, Theorem 5.2] under the restriction $\alpha, \beta \geq -1/2$.

We remark that Wróbel [13, Corollary 4.3] recently proved a multiplier result for Jacobi expansions, which is related to our multiplier results in Section 5. In his proof, Wróbel makes use of the integral representation for the Jacobi-Poisson kernel derived in [7, Proposition 4.1] (which is restricted to $\alpha, \beta \geq -1/2$); see [13, Section 4]. The Jacobi-Poisson kernel formula obtained in Proposition 2.3 should thus make it possible to extend Wróbel’s result to a wider range of α, β than that admitted in [13, Corollary 4.3]; this, however, remains to be investigated.

The paper is organized as follows. In Section 2 we derive an integral representation of the Jacobi-Poisson kernel valid for all $\alpha, \beta > -1$. Section 3 contains various facts and preparatory results needed for kernel estimates. In Section 4 we prove standard estimates for kernels associated with the operators mentioned above. This leads to our main results in Section 5, saying

that the operators in question can be interpreted as Calderón-Zygmund operators and giving, as a consequence, their L^p mapping properties. Finally, Section 6 is devoted to sharp estimates of the Jacobi-Poisson kernel.

Throughout the paper we use a fairly standard notation with essentially all symbols referring to the space of homogeneous type $([0, \pi], d\mu_{\alpha, \beta}, |\cdot|)$. Since the distance in this space is the Euclidean one, the ball denoted $B(\theta, r)$ is simply the interval $(\theta - r, \theta + r) \cap [0, \pi]$. When writing estimates, we will frequently use the notation $X \lesssim Y$ to indicate that $X \leq CY$ with a positive constant C independent of significant quantities. We shall write $X \simeq Y$ when simultaneously $X \lesssim Y$ and $Y \lesssim X$.

2. JACOBI-POISSON KERNEL

Let $\alpha, \beta > -1$. The Jacobi-Poisson kernel is given by (see [7, Section 2])

$$H_t^{\alpha, \beta}(\theta, \varphi) = \sum_{n=0}^{\infty} e^{-t|n + \frac{\alpha + \beta + 1}{2}|} \mathcal{P}_n^{\alpha, \beta}(\theta) \mathcal{P}_n^{\alpha, \beta}(\varphi);$$

here $t > 0$ and $\theta, \varphi \in [0, \pi]$, and $\mathcal{P}_n^{\alpha, \beta}$ are the classical Jacobi trigonometric polynomials, normalized in $L^2(d\mu_{\alpha, \beta})$. Notice that the fraction $\frac{\alpha + \beta + 1}{2}$ may be negative. Defining the auxiliary kernel

$$\mathbb{H}_t^{\alpha, \beta}(\theta, \varphi) := \sum_{n=0}^{\infty} e^{-t(n + \frac{\alpha + \beta + 1}{2})} \mathcal{P}_n^{\alpha, \beta}(\theta) \mathcal{P}_n^{\alpha, \beta}(\varphi),$$

the Jacobi-Poisson kernel can be written as

$$(1) \quad H_t^{\alpha, \beta}(\theta, \varphi) = \mathbb{H}_t^{\alpha, \beta}(\theta, \varphi) + \chi_{\{\alpha + \beta < -1\}} 2^{\alpha + \beta + 2} c_{\alpha, \beta} \sinh\left(\frac{\alpha + \beta + 1}{2} t\right),$$

where

$$c_{\alpha, \beta} := \frac{\Gamma(\alpha + \beta + 2)}{2^{\alpha + \beta + 1} \Gamma(\alpha + 1) \Gamma(\beta + 1)}.$$

The last term in (1) is nonzero when $\alpha + \beta < -1$. As we shall see later, there are important cancellations between the two terms in (1) for large t .

The kernel $\mathbb{H}_t^{\alpha, \beta}(\theta, \varphi)$ can be computed explicitly by means of Bailey's formula, see [1, p. 385–387]. More precisely, we have

$$(2) \quad \begin{aligned} \mathbb{H}_t^{\alpha, \beta}(\theta, \varphi) &= c_{\alpha, \beta} \frac{\sinh \frac{t}{2}}{\left(\cosh \frac{t}{2}\right)^{\alpha + \beta + 2}} \\ &\times F_4\left(\frac{\alpha + \beta + 2}{2}, \frac{\alpha + \beta + 3}{2}; \alpha + 1, \beta + 1; \left(\frac{\sin \frac{\theta}{2} \sin \frac{\varphi}{2}}{\cosh \frac{t}{2}}\right)^2, \left(\frac{\cos \frac{\theta}{2} \cos \frac{\varphi}{2}}{\cosh \frac{t}{2}}\right)^2\right), \end{aligned}$$

for $t > 0$ and $\theta, \varphi \in [0, \pi]$. Here F_4 is Appel's hypergeometric function of two variables defined by the series

$$F_4(a_1, a_2; b_1, b_2; x, y) = \sum_{m, n=0}^{\infty} \frac{(a_1)_{m+n} (a_2)_{m+n}}{(b_1)_m (b_2)_n m! n!} x^m y^n,$$

where $(a)_n$ means the Pochhammer symbol, $(a)_n = a(a+1) \cdots (a+n-1)$ for $n \geq 1$ and $(a)_0 = 1$. This double power series is known to converge absolutely when $\sqrt{|x|} + \sqrt{|y|} < 1$, cf. [3, Chapter V, Section 5.7.2]. From this expression, the positivity of $\mathbb{H}_t^{\alpha, \beta}(\theta, \varphi)$ can easily be seen. Moreover, (2) provides a holomorphic extension of $\mathbb{H}_t^{\alpha, \beta}(\theta, \varphi)$ as a function of the

parameters $\alpha, \beta > -1$ to the region $\{(\alpha, \beta) \in \mathbb{C}^2 : \Re \alpha, \Re \beta > -1\}$. Indeed, with $t > 0$ and $\theta, \varphi \in [0, \pi]$ fixed, the hypergeometric series in (2) is a sum of holomorphic functions of (α, β) converging locally uniformly in the region in question (the latter fact can be justified by means of elementary estimates for the Pochhammer symbol). However, the formula (2) does not seem to be convenient from the point of view of kernel estimates. Thus we need a more suitable representation.

In [7, Section 4] the first and second authors derived the following integral representation, valid for $\alpha, \beta \geq -1/2$ (notice that under this restriction $H_t^{\alpha, \beta}(\theta, \varphi)$ coincides with $\mathbb{H}_t^{\alpha, \beta}(\theta, \varphi)$):

$$(3) \quad \mathbb{H}_t^{\alpha, \beta}(\theta, \varphi) = c_{\alpha, \beta} \sinh \frac{t}{2} \iint \frac{d\Pi_\alpha(u) d\Pi_\beta(v)}{(\cosh \frac{t}{2} - 1 + q(\theta, \varphi, u, v))^{\alpha + \beta + 2}}, \quad t > 0, \quad \theta, \varphi \in [0, \pi],$$

where

$$q(\theta, \varphi, u, v) = 1 - u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2},$$

and the measure $d\Pi_\alpha$ is defined in the following way. For $\alpha > -1/2$, we let

$$(4) \quad \Pi_\alpha(u) := \frac{\Gamma(\alpha + 1)}{\sqrt{\pi} \Gamma(\alpha + 1/2)} \int_0^u (1 - w^2)^{\alpha - 1/2} dw,$$

which is an odd function in $-1 < u < 1$. Then $d\Pi_\alpha$ is a probability measure in $[-1, 1]$. As $\alpha \rightarrow -1/2$, one finds that $d\Pi_\alpha$ converges weakly to the measure $d\Pi_{-1/2} := \frac{1}{2}(\delta_{-1} + \delta_1)$, where $\delta_{\pm 1}$ denotes a point mass at ± 1 .

Now we observe that (4) can be extended to all complex $\alpha \neq -1/2$ with $\Re \alpha > -1$. Then the (distribution) derivative

$$d\Pi_\alpha(u) = \frac{\Gamma(\alpha + 1)}{\sqrt{\pi} \Gamma(\alpha + 1/2)} (1 - u^2)^{\alpha - 1/2} du$$

is a local complex measure in $(-1, 1)$. For $\alpha \in (-1, -1/2)$ real, its density is negative, even and not integrable in $(-1, 1)$. If ϕ is a continuous function in $(-1, 1)$ and $\phi(u) = \mathcal{O}(1 - u)$ as $u \rightarrow 1$, then the integral $I(\alpha) = \int_0^1 \phi(u) d\Pi_\alpha(u)$ is well defined. As a function of α , this integral is analytic in $\{\alpha : \Re \alpha > -1, \alpha \neq -1/2\}$. Since $|I(\alpha)| \lesssim |\alpha + 1/2| \int_0^1 (1 - u^2)^{\Re \alpha + 1/2} du \rightarrow 0$ as $\alpha \rightarrow -1/2$, we see that $I(\alpha)$ is actually analytic in $\{\alpha : \Re \alpha > -1\}$ and $I(-1/2) = 0$. More generally, if $\phi_{\alpha, \beta}(u)$ is continuous in (u, α, β) and analytic in (α, β) for $-1 < u < 1$ and $\Re \alpha, \Re \beta > -1$, and $\phi_{\alpha, \beta}(u) = \mathcal{O}(1 - u)$ locally uniformly in (α, β) , then $I(\alpha, \beta) = \int_0^1 \phi_{\alpha, \beta}(u) d\Pi_\alpha(u)$ will be analytic in (α, β) in $\Re \alpha, \Re \beta > -1$. Under analogous assumptions, this also extends to functions $\phi_{\alpha, \beta}(u, v)$ and the double integral $I(\alpha, \beta) = \iint_{(0, 1)^2} \phi_{\alpha, \beta}(u, v) d\Pi_\alpha(u) d\Pi_\beta(v)$, if one assumes $\phi_{\alpha, \beta}(u, v) = \mathcal{O}((1 - u)(1 - v))$ locally uniformly in α and β .

The measures $d\Pi_\alpha$ will now be used to extend the representation (3) to the range $\alpha, \beta > -1$. Denote

$$(5) \quad \Psi^{\alpha, \beta}(t, \theta, \varphi, u, v) := \frac{c_{\alpha, \beta} \sinh \frac{t}{2}}{(\cosh \frac{t}{2} - 1 + q(\theta, \varphi, u, v))^{\alpha + \beta + 2}}.$$

Taking the even parts of $\Psi^{\alpha, \beta}(t, \theta, \varphi, u, v)$ in u and v , we also define

$$\Psi_E^{\alpha, \beta}(t, \theta, \varphi, u, v) := \frac{1}{4} \sum_{\xi, \eta = \pm 1} \Psi^{\alpha, \beta}(t, \theta, \varphi, \xi u, \eta v).$$

Notice that by (3) and symmetry reasons, for $\alpha, \beta \geq -1/2$ we have

$$(6) \quad \mathbb{H}_t^{\alpha, \beta}(\theta, \varphi) = 4 \iint_{(0,1]^2} \Psi_E^{\alpha, \beta}(t, \theta, \varphi, u, v) d\Pi_\alpha(u) d\Pi_\beta(v).$$

We can now state a general integral representation of $\mathbb{H}_t^{\alpha, \beta}(\theta, \varphi)$.

Theorem 2.1. *For all $\alpha, \beta > -1$, $t > 0$ and $\theta, \varphi \in [0, \pi]$,*

$$(7) \quad \begin{aligned} \mathbb{H}_t^{\alpha, \beta}(\theta, \varphi) = & 4 \iint_{(0,1]^2} \left(\Psi_E^{\alpha, \beta}(t, \theta, \varphi, u, v) - \Psi_E^{\alpha, \beta}(t, \theta, \varphi, u, 1) - \Psi_E^{\alpha, \beta}(t, \theta, \varphi, 1, v) \right. \\ & \left. + \Psi_E^{\alpha, \beta}(t, \theta, \varphi, 1, 1) \right) d\Pi_\alpha(u) d\Pi_\beta(v) \\ & + 2 \int_{(0,1]} \left(\Psi_E^{\alpha, \beta}(t, \theta, \varphi, u, 1) - \Psi_E^{\alpha, \beta}(t, \theta, \varphi, 1, 1) \right) d\Pi_\alpha(u) \\ & + 2 \int_{(0,1]} \left(\Psi_E^{\alpha, \beta}(t, \theta, \varphi, 1, v) - \Psi_E^{\alpha, \beta}(t, \theta, \varphi, 1, 1) \right) d\Pi_\beta(v) \\ & + \Psi_E^{\alpha, \beta}(t, \theta, \varphi, 1, 1). \end{aligned}$$

Proof. For $\alpha, \beta \geq -1/2$, (7) is an easy consequence of (6). With $\phi_{\alpha, \beta}(u) = \Psi_E^{\alpha, \beta}(t, \theta, \varphi, u, 1) - \Psi_E^{\alpha, \beta}(t, \theta, \varphi, 1, 1)$, the second integral in (7) is of the form $I(\alpha, \beta)$ just described; observe that $\phi_{\alpha, \beta}(u) = \mathcal{O}(1-u)$, since the derivative $\partial \Psi_E^{\alpha, \beta} / \partial u$ is bounded locally uniformly in α and β . The third integral in (7) is similar. For the double integral, we let

$$\phi_{\alpha, \beta}(u, v) = \Psi_E^{\alpha, \beta}(t, \theta, \varphi, u, v) - \Psi_E^{\alpha, \beta}(t, \theta, \varphi, u, 1) - \Psi_E^{\alpha, \beta}(t, \theta, \varphi, 1, v) + \Psi_E^{\alpha, \beta}(t, \theta, \varphi, 1, 1),$$

and get a double integral of type $I(\alpha, \beta)$.

The conclusion is that the right-hand side of (7) is analytic in $(\alpha, \beta) \in \{z : \Re z > -1\}^2$. Theorem 2.1 follows, since the left-hand side is also analytic. \square

We remark that in Theorem 2.1 it does not matter whether one integrates over the open interval $(0, 1)$ or over $(0, 1]$, even when the measure is $d\Pi_{-1/2}$. But in the sequel, it will be more convenient to use $(0, 1]$.

Next we restate the formula of Theorem 2.1 in order to obtain a more suitable representation of $\mathbb{H}_t^{\alpha, \beta}(\theta, \varphi)$ for the kernel estimates in Section 4. Recall that for $-1 < \alpha < -1/2$, $\Pi_\alpha(u)$ is an odd function, which is negative for $u > 0$. It can easily be verified that the density $|\Pi_\alpha(u)|$ defines a finite measure on $[-1, 1]$. In fact, we have the following.

Lemma 2.2. *Let $-1 < \alpha < -1/2$ be fixed. Then*

$$|\Pi_\alpha(u)| \simeq |u|(1-|u|)^{\alpha+1/2} \simeq |u| \frac{d\Pi_{\alpha+1}(u)}{du}, \quad u \in (-1, 1).$$

Proof. These three quantities are even in u , and we need consider only $u \in (0, 1)$. It is enough to observe that then $|\Pi_\alpha(u)| \simeq \int_0^u (1-w)^{\alpha-1/2} dw$. \square

Proposition 2.3. *Let $t > 0$ and $\theta, \varphi \in [0, \pi]$.*

(i) *If $\alpha, \beta \geq -1/2$, then*

$$\mathbb{H}_t^{\alpha, \beta}(\theta, \varphi) = \iint \Psi^{\alpha, \beta}(t, \theta, \varphi, u, v) d\Pi_\alpha(u) d\Pi_\beta(v).$$

(ii) If $-1 < \alpha < -1/2 \leq \beta$, then

$$\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi) = \iint \left\{ -\partial_u \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) \Pi_\alpha(u) du d\Pi_\beta(v) \right. \\ \left. + \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) d\Pi_{-1/2}(u) d\Pi_\beta(v) \right\}.$$

(iii) If $-1 < \beta < -1/2 \leq \alpha$, then

$$\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi) = \iint \left\{ -\partial_v \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) d\Pi_\alpha(u) \Pi_\beta(v) dv \right. \\ \left. + \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) d\Pi_\alpha(u) d\Pi_{-1/2}(v) \right\}.$$

(iv) If $-1 < \alpha, \beta < -1/2$, then

$$\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi) = \iint \left\{ \partial_u \partial_v \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) \Pi_\alpha(u) du \Pi_\beta(v) dv \right. \\ - \partial_u \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) \Pi_\alpha(u) du d\Pi_{-1/2}(v) \\ - \partial_v \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) d\Pi_{-1/2}(u) \Pi_\beta(v) dv \\ \left. + \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) d\Pi_{-1/2}(u) d\Pi_{-1/2}(v) \right\}.$$

Proof. Item (i) is just (3). To prove the remaining items, we combine Theorem 2.1, Lemma 2.2 and symmetries of the quantity $\Psi_E^{\alpha,\beta}(t, \theta, \varphi, u, v)$, its derivatives in u and v , and the measures involved. We give further details in case of (ii), leaving similar proofs of (iii) and (iv) to the reader.

Assume that $-1 < \alpha < -1/2 \leq \beta$. Since $d\Pi_\beta$ is a symmetric probability measure on $[-1, 1]$ and has no atom at 0, formula (7) reduces to

$$\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi) = 4 \iint_{(0,1]^2} \left(\Psi_E^{\alpha,\beta}(t, \theta, \varphi, u, v) - \Psi_E^{\alpha,\beta}(t, \theta, \varphi, 1, v) \right) d\Pi_\alpha(u) d\Pi_\beta(v) \\ + 2 \int_{(0,1]} \Psi_E^{\alpha,\beta}(t, \theta, \varphi, 1, v) d\Pi_\beta(v) \\ \equiv I_1 + I_2.$$

Then, expressing $\Psi_E^{\alpha,\beta}$ via $\Psi^{\alpha,\beta}$ and making use of the symmetry of $d\Pi_\beta$, we see that

$$I_2 = 4 \iint_{(0,1]^2} \Psi_E^{\alpha,\beta}(t, \theta, \varphi, u, v) d\Pi_{-1/2}(u) d\Pi_\beta(v) \\ = \iint \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) d\Pi_{-1/2}(u) d\Pi_\beta(v).$$

In I_1 we integrate by parts in the u variable, which is legitimate in view of Lemma 2.2. Observe that the integrand in I_1 vanishes for $u = 1$ and that $\Pi_\alpha(0) = 0$. We get

$$I_1 = -4 \iint_{(0,1]^2} \partial_u \Psi_E^{\alpha,\beta}(t, \theta, \varphi, u, v) \Pi_\alpha(u) du d\Pi_\beta(v).$$

Inserting the definition of the symmetrization $\Psi_E^{\alpha,\beta}$, one easily finds that

$$I_1 = - \iint \partial_u \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) \Pi_\alpha(u) du d\Pi_\beta(v).$$

The conclusion follows. \square

Remark 2.4. All the representations of $\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi)$ contained in Proposition 2.3 are positive in the sense that each of the double integrals (there are one of these in (i), two in (ii) and in (iii), and four in (iv)) is nonnegative.

3. PREPARATORY RESULTS

In this section we gather various technical results, altogether forming a transparent and convenient method of proving standard estimates for kernels defined via the Jacobi-Poisson kernel. The essence of this technique is a uniform way of handling double integrals against products of measures of type $d\Pi_\gamma$ and $\Pi_\gamma(u) du$, independently of the integrands. Then the expressions one has to estimate contain only elementary functions and are relatively simple.

The result below, which is a generalization of [7, Lemma 4.3], plays a crucial role in our method to prove kernel estimates. It provides a link from estimates emerging from the integral representation of $\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi)$, see Proposition 2.3, to the standard estimates related to the space of homogeneous type $([0, \pi], d\mu_{\alpha,\beta}, |\cdot|)$.

Lemma 3.1. Let $\alpha, \beta > -1$. Assume that $\xi_1, \xi_2, \kappa_1, \kappa_2 \geq 0$ are fixed and such that $\alpha + \xi_1 + \kappa_1, \beta + \xi_2 + \kappa_2 \geq -1/2$. Then, uniformly in $\theta, \varphi \in [0, \pi], \theta \neq \varphi$,

$$\begin{aligned} & \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{2\xi_1} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{2\xi_2} \iint \frac{d\Pi_{\alpha+\xi_1+\kappa_1}(u) d\Pi_{\beta+\xi_2+\kappa_2}(v)}{(q(\theta, \varphi, u, v))^{\alpha+\beta+\xi_1+\xi_2+3/2}} \\ & \lesssim \frac{1}{\mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}, \\ & \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{2\xi_1} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{2\xi_2} \iint \frac{d\Pi_{\alpha+\xi_1+\kappa_1}(u) d\Pi_{\beta+\xi_2+\kappa_2}(v)}{(q(\theta, \varphi, u, v))^{\alpha+\beta+\xi_1+\xi_2+2}} \\ & \lesssim \frac{1}{|\theta - \varphi| \mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}. \end{aligned}$$

Note that for any $\alpha, \beta > -1$ fixed, the $\mu_{\alpha,\beta}$ measure of the interval $B(\theta, |\theta - \varphi|)$ can be described as follows, see [7, Lemma 4.2],

$$(8) \quad \mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|)) \simeq |\theta - \varphi|(\theta + \varphi)^{2\alpha+1}(\pi - \theta + \pi - \varphi)^{2\beta+1}, \quad \theta, \varphi \in [0, \pi].$$

Notice also that the right-hand sides of the estimates in Lemma 3.1 are always larger than a positive constant, uniformly in $\theta, \varphi \in [0, \pi]$. This fact will be used in the sequel without further mention.

To prove Lemma 3.1, we need item (b) in the lemma below. This is a generalization of [7, Lemma 4.4] stated in [9, Lemma 2.3]. Item (a) contains estimates obtained in [8, Lemma 5.3] and will be used in Section 6 below to show sharp estimates of the Jacobi-Poisson kernel.

Lemma 3.2. Let $\kappa \geq 0$ and γ and ν be such that $\gamma > \nu + 1/2 \geq 0$. Then

(a)

$$\int \frac{d\Pi_\nu(s)}{(D - Bs)^\kappa (A - Bs)^\gamma} \simeq \frac{1}{(D - B)^\kappa A^{\nu+1/2} (A - B)^{\gamma-\nu-1/2}}, \quad 0 \leq B < A \leq D;$$

(b)

$$\int \frac{d\Pi_{\nu+\kappa}(s)}{(A - Bs)^\gamma} \lesssim \frac{1}{A^{\nu+1/2} (A - B)^{\gamma-\nu-1/2}}, \quad 0 \leq B < A.$$

Proof. Part (a) is proved in [8]. Part (b) can easily be deduced from (a) since the integral to be estimated is controlled by the same integral with $\kappa = 0$. \square

Proof of Lemma 3.1. The reasoning is a combination of the arguments given in the proofs of [9, Lemma 2.1] and [7, Lemma 4.3]. Since $(\theta - \varphi)^2 \lesssim q(\theta, \varphi, u, v)$, it suffices to verify the first estimate. Further, we may reduce the task to showing that

$$(9) \quad \iint \frac{d\Pi_{\alpha+\kappa_1}(u) d\Pi_{\beta+\kappa_2}(v)}{(q(\theta, \varphi, u, v))^{\alpha+\beta+3/2}} \lesssim \frac{1}{\mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}, \quad \theta, \varphi \in [0, \pi], \quad \theta \neq \varphi,$$

under the assumption $\alpha + \kappa_1, \beta + \kappa_2 \geq -1/2$. Indeed, applying (9) with $\alpha + \xi_1, \beta + \xi_2$ instead of α, β , and then using (8), we obtain

$$\begin{aligned} & \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{2\xi_1} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{2\xi_2} \iint \frac{d\Pi_{\alpha+\xi_1+\kappa_1}(u) d\Pi_{\beta+\xi_2+\kappa_2}(v)}{(q(\theta, \varphi, u, v))^{\alpha+\beta+\xi_1+\xi_2+3/2}} \\ & \lesssim (\theta + \varphi)^{2\xi_1} (\pi - \theta + \pi - \varphi)^{2\xi_2} \frac{1}{\mu_{\alpha+\xi_1, \beta+\xi_2}(B(\theta, |\theta - \varphi|))} \simeq \frac{1}{\mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}. \end{aligned}$$

To prove (9), it is convenient to distinguish two cases.

Case 1: $\alpha, \beta \in (-1, -1/2)$.

Taking into account the estimates, see [7, (21)],

$$|\theta - \varphi|^2 \simeq 2 \sin^2 \frac{\theta - \varphi}{4} \leq q(\theta, \varphi, u, v) \leq 2 \cos^2 \frac{\theta - \varphi}{4} \leq 2, \quad \theta, \varphi \in [0, \pi], \quad u, v \in [-1, 1],$$

and the fact that $d\Pi_{\alpha+\kappa_1}$ and $d\Pi_{\beta+\kappa_2}$ are finite, we get

$$\iint \frac{d\Pi_{\alpha+\kappa_1}(u) d\Pi_{\beta+\kappa_2}(v)}{(q(\theta, \varphi, u, v))^{\alpha+\beta+3/2}} \lesssim \frac{1}{|\theta - \varphi|^{2\alpha+1} |\theta - \varphi|^{2\beta+1} |\theta - \varphi|} + \chi_{\{\alpha+\beta+3/2 < 0\}}.$$

Then using the inequalities $|\theta - \varphi| \leq \theta + \varphi$ and $|\theta - \varphi| \leq \pi - \theta + \pi - \varphi$ together with (8), we obtain (9).

Case 2: At least one of the parameters α, β is in $[-1/2, \infty)$, say $\beta \geq -1/2$.

Proceeding as in the proof of [7, Lemma 4.3] but applying Lemma 3.2 (b) instead of [7, Lemma 4.4] to the integral against $d\Pi_{\beta+\kappa_2}$, we see that

$$\iint \frac{d\Pi_{\alpha+\kappa_1}(u) d\Pi_{\beta+\kappa_2}(v)}{(q(\theta, \varphi, u, v))^{\alpha+\beta+3/2}} \lesssim \frac{1}{(\pi - \theta + \pi - \varphi)^{2\beta+1}} \int \frac{d\Pi_{\alpha+\kappa_1}(u)}{(q(\theta, \varphi, u, 1))^{\alpha+1}}.$$

When $\alpha \geq -1/2$, another application of Lemma 3.2 (b) leads to (9), see the proof of [7, Lemma 4.3]. If $\alpha \in (-1, -1/2)$ we can apply the arguments from Case 1 getting

$$\int \frac{d\Pi_{\alpha+\kappa_1}(u)}{(q(\theta, \varphi, u, 1))^{\alpha+1}} \lesssim \frac{1}{|\theta - \varphi|^{2\alpha+2}} \leq \frac{1}{(\theta + \varphi)^{2\alpha+1} |\theta - \varphi|}.$$

Using now (8), we arrive at the desired conclusion.

The proof of Lemma 3.1 is complete. \square

The remaining part of this section embraces various technical results, which will allow us to control the relevant kernels by means of the estimates from Lemma 3.1. To state the next lemma and also for further use, we introduce the following notation. We will omit the arguments and

write briefly \mathfrak{q} instead of $q(\theta, \varphi, u, v)$, when it does not lead to confusion. For a given parameter $\lambda \in \mathbb{R}$, we define the auxiliary function

$$\Psi^\lambda(t, \mathfrak{q}) := \frac{\sinh \frac{t}{2}}{(\cosh \frac{t}{2} - 1 + \mathfrak{q})^\lambda},$$

so that $\Psi^{\alpha, \beta}(t, \theta, \varphi, u, v) = c_{\alpha, \beta} \Psi^{\alpha + \beta + 2}(t, \mathfrak{q})$; see (5).

Lemma 3.3. *Let $\lambda \in \mathbb{R}$, $M, N \in \mathbb{N} = \{0, 1, 2, \dots\}$ and $K, R, L \in \{0, 1\}$ be fixed. Then*

$$\begin{aligned} & |\partial_u^K \partial_v^R \partial_\varphi^L \partial_\theta^N \partial_t^M \Psi^\lambda(t, \mathfrak{q})| \\ & \lesssim \sum_{k, r=0, 1, 2} \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{Kk} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{Rr} \frac{1}{(t^2 + \mathfrak{q})^{\lambda + (L + N + M - 1 + Kk + Rr)/2}}, \end{aligned}$$

uniformly in $t \in (0, 1]$, $\theta, \varphi \in [0, \pi]$ and $u, v \in [-1, 1]$.

To prove this lemma, we need two preparatory results. One of them is Faà di Bruno's formula for the N th derivative, $N \geq 1$, of the composition of two functions (see [5] for the related references and interesting historical remarks). With D denoting the ordinary derivative, it reads

$$(10) \quad D^N(g \circ f)(\theta) = \sum \frac{N!}{j_1! \cdots j_N!} (D^{j_1 + \dots + j_N} g) \circ f(\theta) \left(\frac{D^1 f(\theta)}{1!} \right)^{j_1} \cdots \left(\frac{D^N f(\theta)}{N!} \right)^{j_N},$$

where the summation runs over all $j_1, \dots, j_N \geq 0$ such that $j_1 + 2j_2 + \dots + Nj_N = N$. Further, in the proof of Lemma 3.3 we will make use of the following bounds given in [7].

Lemma 3.4 ([7, Lemma 4.5]). *For all $\theta, \varphi \in [0, \pi]$ and $u, v \in [-1, 1]$, one has*

$$|\partial_\theta \mathfrak{q}| \lesssim \sqrt{\mathfrak{q}} \quad \text{and} \quad |\partial_\varphi \mathfrak{q}| \lesssim \sqrt{\mathfrak{q}}.$$

Proof of Lemma 3.3. Given $\lambda \in \mathbb{R}$, we introduce the auxiliary function

$$\tilde{\Psi}^\lambda(t, \mathfrak{q}) := \frac{1}{\sinh \frac{t}{2}} \Psi^\lambda(t, \mathfrak{q}) = \frac{1}{(\cosh \frac{t}{2} - 1 + \mathfrak{q})^\lambda}.$$

We first reduce our task to showing the estimate

$$(11) \quad \begin{aligned} & |\partial_u^K \partial_v^R \partial_\varphi^L \partial_\theta^N \tilde{\Psi}^\lambda(t, \mathfrak{q})| \\ & \lesssim \sum_{k, r=0, 1, 2} \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{Kk} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{Rr} \frac{1}{(t^2 + \mathfrak{q})^{\lambda + (L + N + Kk + Rr)/2}} \end{aligned}$$

for $t \in (0, 1]$, $\theta, \varphi \in [0, \pi]$ and $u, v \in [-1, 1]$; here $\lambda \in \mathbb{R}$, $N \in \mathbb{N}$ and $K, R, L \in \{0, 1\}$ are fixed.

Observe that

$$\Psi^\lambda(t, \mathfrak{q}) = c_\lambda \begin{cases} \partial_t \frac{1}{(\cosh \frac{t}{2} - 1 + \mathfrak{q})^{\lambda-1}}, & \lambda \neq 1 \\ \partial_t \log(\cosh \frac{t}{2} - 1 + \mathfrak{q}), & \lambda = 1 \end{cases},$$

where c_λ is a constant, possibly negative. Using Faà di Bruno's formula (10) with $f(t) = \cosh \frac{t}{2} - 1 + \mathfrak{q}$ and either $g(x) = x^{-\lambda+1}$ or $g(x) = \log x$, we obtain

$$\begin{aligned} \partial_t^M \Psi^\lambda(t, \mathfrak{q}) &= c_\lambda \partial_t^{M+1} (g \circ f)(t) \\ &= \sum_{\substack{j_i \geq 0 \\ j_1 + \dots + (M+1)j_{M+1} = M+1}} C_{\lambda, j} \left(\sinh \frac{t}{2} \right)^{\sum_{\text{odd } i} j_i} \left(\cosh \frac{t}{2} \right)^{\sum_{\text{even } i} j_i} \tilde{\Psi}^{\lambda-1 + \sum_i j_i}(t, \mathfrak{q}), \end{aligned}$$

where the $C_{\lambda,j}$ are constants, possibly zero. Differentiating these identities with respect to θ, φ, u, v and then applying (11) and the relations

$$\cosh \frac{t}{2} \simeq 1, \quad \sinh \frac{t}{2} \simeq t \leq \sqrt{t^2 + \mathbf{q}}, \quad t \in (0, 1],$$

we see that

$$|\partial_u^K \partial_v^R \partial_\varphi^L \partial_\theta^N \partial_t^M \Psi^\lambda(t, \mathbf{q})| \lesssim \sum_{\substack{j_i \geq 0 \\ j_1 + \dots + (M+1)j_{M+1} = M+1}} \sum_{k,r=0,1,2} \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{Kk} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{Rr} \\ \times \frac{1}{(t^2 + \mathbf{q})^{\lambda-1+\sum_i j_i - (\sum_{\text{odd } i} j_i)/2 + (L+N+Kk+Rr)/2}}.$$

Now by the boundedness of \mathbf{q} and the inequality

$$(12) \quad \sum_i j_i - \frac{1}{2} \sum_{\text{odd } i} j_i \leq \frac{M+1}{2},$$

forced by the constraint $j_1 + \dots + (M+1)j_{M+1} = M+1$, we get the asserted estimate. Thus it remains to prove (11).

We assume that $N \geq 1$. The simpler case $N = 0$ is left to the reader. Taking into account the relations

$$\partial_\theta^{2m} \mathbf{q} = (-4)^{-m} (\mathbf{q} - 1), \quad \partial_\theta^{2m-1} \mathbf{q} = (-4)^{1-m} \partial_\theta \mathbf{q}, \quad m \geq 1,$$

see [7, Section 4], and using Faà di Bruno's formula with $f(\theta) = \cosh \frac{t}{2} - 1 + \mathbf{q}$ and $g(x) = x^{-\lambda}$, we get

$$\partial_\theta^N \tilde{\Psi}^\lambda(t, \mathbf{q}) = \sum_{\substack{j_i \geq 0 \\ j_1 + \dots + Nj_N = N}} c_{\lambda,j} \frac{1}{(\cosh \frac{t}{2} - 1 + \mathbf{q})^{\lambda + \sum_i j_i}} (\mathbf{q} - 1)^{\sum_{\text{even } i} j_i} (\partial_\theta \mathbf{q})^{\sum_{\text{odd } i} j_i},$$

where $c_{\lambda,j}$ are constants. Further, keeping in mind that $L, R, K \in \{0, 1\}$ and applying repeatedly Leibniz' rule, we see that $\partial_\varphi^L \partial_\theta^N \tilde{\Psi}^\lambda(t, \mathbf{q})$ is a sum of terms of the form constant times

$$\frac{1}{(\cosh \frac{t}{2} - 1 + \mathbf{q})^{\lambda + \sum_i j_i + l_1}} (\partial_\varphi \mathbf{q})^{l_1 + l_2} (\mathbf{q} - 1)^{\sum_{\text{even } i} j_i - l_2} (\partial_\theta \mathbf{q})^{\sum_{\text{odd } i} j_i - l_3} (\partial_\varphi \partial_\theta \mathbf{q})^{l_3},$$

where the indices run over the set described by the conditions $j_i \geq 0$, $j_1 + \dots + Nj_N = N$, $l_1, l_2, l_3 \geq 0$, $l_1 + l_2 + l_3 = L$ and the exponents of $\mathbf{q} - 1$ and $\partial_\theta \mathbf{q}$ are nonnegative. Similarly, $\partial_v^R \partial_\varphi^L \partial_\theta^N \tilde{\Psi}^\lambda(t, \mathbf{q})$ is a sum of terms of the form constant times

$$\frac{1}{(\cosh \frac{t}{2} - 1 + \mathbf{q})^{\lambda + \sum_i j_i + l_1 + r_1}} (\partial_v \mathbf{q})^{r_1 + r_3} (\partial_\varphi \mathbf{q})^{l_1 + l_2 - r_2} (\partial_v \partial_\varphi \mathbf{q})^{r_2} (\mathbf{q} - 1)^{\sum_{\text{even } i} j_i - l_2 - r_3} \\ \times (\partial_\theta \mathbf{q})^{\sum_{\text{odd } i} j_i - l_3 - r_4} (\partial_v \partial_\theta \mathbf{q})^{r_4} (\partial_\varphi \partial_\theta \mathbf{q})^{l_3 - r_5} (\partial_v \partial_\varphi \partial_\theta \mathbf{q})^{r_5},$$

where also $r_1, \dots, r_5 \geq 0$, $r_1 + \dots + r_5 = R$, $l_1 + l_2 \geq r_2$, $l_3 \geq r_5$. Finally, since the derivative $\partial_u \partial_v \mathbf{q}$ vanishes, $\partial_u^K \partial_v^R \partial_\varphi^L \partial_\theta^N \tilde{\Psi}^\lambda(t, \mathbf{q})$ is a sum of terms of the form constant times

$$\frac{1}{(\cosh \frac{t}{2} - 1 + \mathbf{q})^{\lambda + \sum_i j_i + l_1 + r_1 + k_1}} (\partial_u \mathbf{q})^{k_1 + k_3} (\partial_v \mathbf{q})^{r_1 + r_3} (\partial_\varphi \mathbf{q})^{l_1 + l_2 - r_2 - k_2} (\partial_u \partial_\varphi \mathbf{q})^{k_2} (\partial_v \partial_\varphi \mathbf{q})^{r_2} \\ \times (\mathbf{q} - 1)^{\sum_{\text{even } i} j_i - l_2 - r_3 - k_3} (\partial_\theta \mathbf{q})^{\sum_{\text{odd } i} j_i - l_3 - r_4 - k_4} (\partial_u \partial_\theta \mathbf{q})^{k_4} (\partial_v \partial_\theta \mathbf{q})^{r_4} (\partial_\varphi \partial_\theta \mathbf{q})^{l_3 - r_5 - k_5} \\ \times (\partial_u \partial_\varphi \partial_\theta \mathbf{q})^{k_5} (\partial_v \partial_\varphi \partial_\theta \mathbf{q})^{r_5}.$$

Here we must add the conditions $k_1, \dots, k_5 \geq 0$, $k_1 + \dots + k_5 = K$ and replace $l_1 + l_2 \geq r_2$, $l_3 \geq r_5$ by $l_1 + l_2 \geq r_2 + k_2$, $l_3 \geq r_5 + k_5$. We shall estimate all the factors in this product from above. Since $t \leq 1$, we can replace $\cosh \frac{t}{2} - 1 + \mathfrak{q}$ by $t^2 + \mathfrak{q}$. The quantities \mathfrak{q} and $\partial_\varphi \partial_\theta \mathfrak{q}$ are bounded. Further, we apply Lemma 3.4 to get

$$|\partial_\varphi \mathfrak{q}| + |\partial_\theta \mathfrak{q}| \lesssim \sqrt{\mathfrak{q}} \leq \sqrt{t^2 + \mathfrak{q}}.$$

To deal with the resulting exponent of $1/(t^2 + \mathfrak{q})$, we observe that

$$l_1 - l_2 + l_3 \leq L, \quad \sum_i j_i - \frac{1}{2} \sum_{\text{odd } i} j_i \leq \frac{N}{2},$$

cf. (12). Using also the estimates

$$\begin{aligned} |\partial_u \mathfrak{q}| &\leq \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^2, & |\partial_v \mathfrak{q}| &\leq \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^2, \\ |\partial_\theta \partial_u \mathfrak{q}| + |\partial_\varphi \partial_u \mathfrak{q}| &\leq \sin \frac{\theta}{2} + \sin \frac{\varphi}{2}, & |\partial_\theta \partial_v \mathfrak{q}| + |\partial_\varphi \partial_v \mathfrak{q}| &\leq \cos \frac{\theta}{2} + \cos \frac{\varphi}{2}, \\ |\partial_\varphi \partial_\theta \partial_u \mathfrak{q}| &\leq 1, & |\partial_\varphi \partial_\theta \partial_v \mathfrak{q}| &\leq 1, \end{aligned}$$

we infer that

$$\begin{aligned} |\partial_u^K \partial_v^R \partial_\varphi^L \partial_\theta^N \tilde{\Psi}^\lambda(t, \mathfrak{q})| &\lesssim \sum_{\substack{r_1 + \dots + r_5 = R \\ k_1 + \dots + k_5 = K}} \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{2k_1 + 2k_3 + k_2 + k_4} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{2r_1 + 2r_3 + r_2 + r_4} \\ &\quad \times \frac{1}{(t^2 + \mathfrak{q})^{\lambda + (N + L + 2k_1 + k_2 + k_4 + 2r_1 + r_2 + r_4)/2}}. \end{aligned}$$

Notice that $2k_1 + k_2 + k_4 \in \{0, K, 2K\}$, and similarly $2r_1 + r_2 + r_4 \in \{0, R, 2R\}$. This observation leads directly to (11). The proof of Lemma 3.3 is complete. \square

Define

$$d\Pi_{\alpha, K} = \begin{cases} d\Pi_{-1/2}, & K = 0 \\ d\Pi_{\alpha+1}, & K = 1 \end{cases},$$

and similary for $d\Pi_{\beta, R}$.

Corollary 3.5. *Let $M, N \in \mathbb{N}$ and $L \in \{0, 1\}$ be fixed. The following estimates hold uniformly in $t \in (0, 1]$ and $\theta, \varphi \in [0, \pi]$.*

(i) *If $\alpha, \beta \geq -1/2$, then*

$$|\partial_\varphi^L \partial_\theta^N \partial_t^M H_t^{\alpha, \beta}(\theta, \varphi)| \lesssim \iint \frac{d\Pi_\alpha(u) d\Pi_\beta(v)}{(t^2 + \mathfrak{q})^{\alpha + \beta + 3/2 + (L + N + M)/2}}.$$

(ii) *If $-1 < \alpha < -1/2 \leq \beta$, then*

$$\begin{aligned} |\partial_\varphi^L \partial_\theta^N \partial_t^M H_t^{\alpha, \beta}(\theta, \varphi)| &\lesssim 1 + \sum_{K=0,1} \sum_{k=0,1,2} \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{Kk} \\ &\quad \times \iint \frac{d\Pi_{\alpha, K}(u) d\Pi_\beta(v)}{(t^2 + \mathfrak{q})^{\alpha + \beta + 3/2 + (L + N + M + Kk)/2}}. \end{aligned}$$

(iii) *If $-1 < \beta < -1/2 \leq \alpha$, then*

$$|\partial_\varphi^L \partial_\theta^N \partial_t^M H_t^{\alpha, \beta}(\theta, \varphi)| \lesssim 1 + \sum_{R=0,1} \sum_{r=0,1,2} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{Rr}$$

$$\times \iint \frac{d\Pi_\alpha(u) d\Pi_{\beta,R}(v)}{(t^2 + \mathfrak{q})^{\alpha+\beta+3/2+(L+N+M+Rr)/2}}.$$

(iv) If $-1 < \alpha, \beta < -1/2$, then

$$\begin{aligned} |\partial_\varphi^L \partial_\theta^N \partial_t^M H_t^{\alpha,\beta}(\theta, \varphi)| &\lesssim 1 + \sum_{K,R=0,1} \sum_{k,r=0,1,2} \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{Kk} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{Rr} \\ &\times \iint \frac{d\Pi_{\alpha,K}(u) d\Pi_{\beta,R}(v)}{(t^2 + \mathfrak{q})^{\alpha+\beta+3/2+(L+N+M+Kk+Rr)/2}}. \end{aligned}$$

Proof. All the bounds are direct consequences of the equality (1), Proposition 2.3, Lemma 2.2 and the estimate from Lemma 3.3 (specified to $\lambda = \alpha + \beta + 2$). Here passing with the differentiation in t, θ or φ under integrals against $d\Pi_\gamma$, $\gamma \geq -1/2$, or $\Pi_\gamma(u) du$, $-1 < \gamma < -1/2$, is legitimate and can easily be justified with the aid of Lemma 3.3 and the dominated convergence theorem. \square

Lemma 3.6. *Let $\gamma \in \mathbb{R}$ and $\eta \geq 0$ be fixed. Then*

$$\int_0^1 \frac{t^\eta dt}{(t^2 + \rho)^{\gamma+\eta/2+1/2}} \lesssim \begin{cases} \rho^{-\gamma}, & \gamma > 0 \\ \log(1 + \rho^{-1/2}), & \gamma = 0 \\ 1, & \gamma < 0 \end{cases},$$

uniformly in $0 < \rho \leq 2$.

Proof. This is elementary. For $\gamma = 0$, one has

$$\int_0^1 \frac{t^\eta dt}{(t^2 + \rho)^{\eta/2+1/2}} \leq \int_0^1 \frac{dt}{(t^2 + \rho)^{1/2}} \simeq \int_0^1 \frac{dt}{t + \rho^{1/2}} = \log(1 + \rho^{-1/2}).$$

\square

The next lemma will be frequently used in Section 4 to prove the relevant kernel estimates. Only the cases $p \in \{1, 2, \infty\}$ will be needed for our purposes. Other values of p are also of interest, but in connection with operators not considered in this paper.

Lemma 3.7. *Let $K, R \in \{0, 1\}$, $k, r \in \{0, 1, 2\}$, $W \geq 1$, $s \geq 0$ and $1 \leq p \leq \infty$ be fixed. Consider a function $\Upsilon_s^{\alpha,\beta}(t, \theta, \varphi)$ defined on $(0, 1) \times [0, \pi] \times [0, \pi]$ in the following way.*

(i) For $\alpha, \beta \geq -1/2$,

$$\Upsilon_s^{\alpha,\beta}(t, \theta, \varphi) := \iint \frac{d\Pi_\alpha(u) d\Pi_\beta(v)}{(t^2 + \mathfrak{q})^{\alpha+\beta+3/2+W/(2p)+s/2}}.$$

(ii) For $-1 < \alpha < -1/2 \leq \beta$,

$$\Upsilon_s^{\alpha,\beta}(t, \theta, \varphi) := \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{Kk} \iint \frac{d\Pi_{\alpha,K}(u) d\Pi_\beta(v)}{(t^2 + \mathfrak{q})^{\alpha+\beta+3/2+W/(2p)+Kk/2+s/2}}.$$

(iii) For $-1 < \beta < -1/2 \leq \alpha$,

$$\Upsilon_s^{\alpha,\beta}(t, \theta, \varphi) := \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{Rr} \iint \frac{d\Pi_\alpha(u) d\Pi_{\beta,R}(v)}{(t^2 + \mathfrak{q})^{\alpha+\beta+3/2+W/(2p)+Rr/2+s/2}}.$$

(iv) For $-1 < \alpha, \beta < -1/2$,

$$\begin{aligned} \Upsilon_s^{\alpha, \beta}(t, \theta, \varphi) &:= \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{Kk} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{Rr} \\ &\quad \times \iint \frac{d\Pi_{\alpha, K}(u) d\Pi_{\beta, R}(v)}{(t^2 + \mathfrak{q})^{\alpha + \beta + 3/2 + W/(2p) + Kk/2 + Rr/2 + s/2}}. \end{aligned}$$

Then the estimate

$$\|1 + \Upsilon_s^{\alpha, \beta}(t, \theta, \varphi)\|_{L^p((0,1), t^{W-1} dt)} \lesssim \frac{1}{|\theta - \varphi|^s} \frac{1}{\mu_{\alpha, \beta}(B(\theta, |\theta - \varphi|))}$$

holds uniformly in $\theta, \varphi \in [0, \pi]$, $\theta \neq \varphi$.

Proof. It is enough to prove the desired estimate without the term 1 in the left-hand side. Further, since $|\theta - \varphi|^2 \lesssim \mathfrak{q}$, it suffices to consider the case $s = 0$. We prove the estimate when $-1 < \alpha, \beta < -1/2$. The remaining cases are left to the reader; they are simpler since then $\alpha + \beta + 3/2 > 0$ and one needs Lemma 3.6 only with $\gamma > 0$.

We first assume that $p < \infty$. Using Minkowski's integral inequality and then Lemma 3.6 with $\gamma = p(\alpha + \beta + 3/2 + Kk/2 + Rr/2)$, $\eta = W - 1$ and $\rho = \mathfrak{q}$, we obtain

$$\begin{aligned} &\|\Upsilon_0^{\alpha, \beta}(t, \theta, \varphi)\|_{L^p((0,1), t^{W-1} dt)} \\ &\leq \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{Kk} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{Rr} \iint \left(\int_0^1 \frac{t^{W-1} dt}{(t^2 + \mathfrak{q})^{p(\alpha + \beta + 3/2 + W/(2p) + Kk/2 + Rr/2)}} \right)^{1/p} \\ &\quad \times d\Pi_{\alpha, K}(u) d\Pi_{\beta, R}(v) \\ &\lesssim \left(\sin \frac{\theta}{2} + \sin \frac{\varphi}{2} \right)^{Kk} \left(\cos \frac{\theta}{2} + \cos \frac{\varphi}{2} \right)^{Rr} \iint \left[\left(\frac{1}{\mathfrak{q}} \right)^{\alpha + \beta + 3/2 + Kk/2 + Rr/2} + 1 \right. \\ &\quad \left. + \left(\log(1 + \mathfrak{q}^{-1/2}) \right)^{1/p} \right] d\Pi_{\alpha, K}(u) d\Pi_{\beta, R}(v). \end{aligned}$$

Now an application of Lemma 3.1 (specified to $\xi_1 = Kk/2$, $\kappa_1 = -\alpha - 1/2$ if $K = 0$ and $\kappa_1 = 1 - k/2$ if $K = 1$, $\xi_2 = Rr/2$, $\kappa_2 = -\beta - 1/2$ if $R = 0$ and $\kappa_2 = 1 - r/2$ if $R = 1$) gives the desired estimate for the expression emerging from the first term in the last integral. As for the remaining two expressions, we observe that $1 \lesssim \log(1 + \mathfrak{q}^{-1/2}) \lesssim \log(1 + |\theta - \varphi|^{-1})$. Moreover, as can be seen from (8), there exists an $\varepsilon = \varepsilon(\alpha, \beta) > 0$ such that

$$\mu_{\alpha, \beta}(B(\theta, |\theta - \varphi|)) \lesssim |\theta - \varphi|^\varepsilon, \quad \theta, \varphi \in [0, \pi].$$

Since the measures $d\Pi_{\alpha, K}$ and $d\Pi_{\beta, R}$ are finite, the conclusion follows.

The case $p = \infty$ can be justified in a similar way by using in the reasoning above the estimate

$$\frac{1}{(t^2 + \mathfrak{q})^{\alpha + \beta + 3/2 + Kk/2 + Rr/2}} \lesssim \left(\frac{1}{\mathfrak{q}} \right)^{\alpha + \beta + 3/2 + Kk/2 + Rr/2} + 1, \quad t \in (0, 1),$$

instead of Lemma 3.6. \square

The next lemma and corollaries are long-time counterparts of Corollary 3.5 and Lemma 3.7.

Lemma 3.8. Assume that $M, N \in \mathbb{N}$ and $L \in \{0, 1\}$ are fixed. Given $\alpha, \beta > -1$, there exists an $\epsilon = \epsilon(\alpha, \beta) > 0$ such that

$$|\partial_\varphi^L \partial_\theta^N \partial_t^M H_t^{\alpha, \beta}(\theta, \varphi)|$$

$$\lesssim e^{-t(|\frac{\alpha+\beta+1}{2}|+\epsilon)} + \chi_{\{N=L=0, \alpha+\beta+1 \neq 0\}} e^{-t|\frac{\alpha+\beta+1}{2}|} + \chi_{\{M=N=L=0, \alpha+\beta+1=0\}},$$

uniformly in $t \geq 1$ and $\theta, \varphi \in [0, \pi]$. Moreover, one can take $\epsilon = (\alpha + \beta + 2) \wedge 1$.

To prove this, it is more convenient to employ the series representation of $H_t^{\alpha, \beta}(\theta, \varphi)$ rather than the formulas from Proposition 2.3.

Proof of Lemma 3.8. For $\alpha, \beta > -1$, $t > 0$ and $\theta, \varphi \in [0, \pi]$ we have

$$(13) \quad H_t^{\alpha, \beta}(\theta, \varphi) = \frac{1}{\mu_{\alpha, \beta}([0, \pi])} e^{-t|\frac{\alpha+\beta+1}{2}|} + \sum_{n=1}^{\infty} e^{-t(n+\frac{\alpha+\beta+1}{2})} \mathcal{P}_n^{\alpha, \beta}(\theta) \mathcal{P}_n^{\alpha, \beta}(\varphi).$$

Denote the sum in (13) by S . To estimate S and its derivatives, we will need suitable bounds for $\partial_{\theta}^N \mathcal{P}_n^{\alpha, \beta}(\theta)$, $N \geq 0$. It is known (see [12, (7.32.2)]) that

$$(14) \quad |\mathcal{P}_n^{\alpha, \beta}(\theta)| \lesssim n^{\alpha+\beta+2}, \quad \theta \in [0, \pi], \quad n \geq 1.$$

Combining this with the identity (cf. [12, (4.21.7)])

$$\partial_{\theta} \mathcal{P}_n^{\alpha, \beta}(\theta) = -\frac{1}{2} \sqrt{n(n+\alpha+\beta+1)} \sin \theta \mathcal{P}_{n-1}^{\alpha+1, \beta+1}(\theta), \quad n \geq 1,$$

we see that for each $N \geq 0$

$$|\partial_{\theta}^N \mathcal{P}_n^{\alpha, \beta}(\theta)| \lesssim n^{3N+\alpha+\beta+2}, \quad \theta \in [0, \pi], \quad n \geq 1.$$

In view of these facts, the series in (13) can be repeatedly differentiated term by term in t, θ and φ , and we get the bounds

$$\begin{aligned} \partial_{\varphi}^L \partial_{\theta}^N \partial_t^M S &\lesssim \sum_{n=1}^{\infty} e^{-t(n+\frac{\alpha+\beta+1}{2})} n^{M+3N+3L+2\alpha+2\beta+4} \\ &= e^{-t(|\frac{\alpha+\beta+1}{2}|+(\alpha+\beta+2)\wedge 1)} \sum_{n=1}^{\infty} e^{-t(n-1)} n^{M+3N+3L+2\alpha+2\beta+4} \\ &\lesssim e^{-t(|\frac{\alpha+\beta+1}{2}|+(\alpha+\beta+2)\wedge 1)}, \end{aligned}$$

uniformly in $t \geq 1$ and $\theta, \varphi \in [0, \pi]$.

Since the other term in (13) is trivial to handle, the conclusion follows. \square

Corollary 3.9. *Let $\alpha, \beta > -1$, $M, N \in \mathbb{N}$, $L \in \{0, 1\}$, $W \geq 1$ and $1 \leq p \leq \infty$ be fixed. Then*

$$\left\| \sup_{\theta, \varphi \in [0, \pi]} |\partial_{\varphi}^L \partial_{\theta}^N \partial_t^M H_t^{\alpha, \beta}(\theta, \varphi)| \right\|_{L^p((1, \infty), t^{W-1} dt)} < \infty,$$

excluding the cases when simultaneously $\alpha + \beta + 1 = 0$ and $M = N = L = 0$ and $p < \infty$.

A strengthened special case of Corollary 3.9 will be needed when we estimate kernels associated with multipliers of Laplace-Stieltjes type.

Corollary 3.10. *Let $\alpha, \beta > -1$ and $L, N \in \{0, 1\}$ be fixed. Then*

$$\left\| e^{t|\frac{\alpha+\beta+1}{2}|} \sup_{\theta, \varphi \in [0, \pi]} |\partial_{\varphi}^L \partial_{\theta}^N H_t^{\alpha, \beta}(\theta, \varphi)| \right\|_{L^{\infty}((1, \infty), dt)} < \infty.$$

4. KERNEL ESTIMATES

Let \mathbb{B} be a Banach space and let $K(\theta, \varphi)$ be a kernel defined on $[0, \pi] \times [0, \pi] \setminus \{(\theta, \varphi) : \theta = \varphi\}$ and taking values in \mathbb{B} . We say that $K(\theta, \varphi)$ is a *standard kernel* in the sense of the space of homogeneous type $([0, \pi], d\mu_{\alpha, \beta}, |\cdot|)$ if it satisfies the so-called *standard estimates*, i.e., the growth estimate

$$(15) \quad \|K(\theta, \varphi)\|_{\mathbb{B}} \lesssim \frac{1}{\mu_{\alpha, \beta}(B(\theta, |\theta - \varphi|))}$$

and the smoothness estimates

$$(16) \quad \|K(\theta, \varphi) - K(\theta', \varphi)\|_{\mathbb{B}} \lesssim \frac{|\theta - \theta'|}{|\theta - \varphi|} \frac{1}{\mu_{\alpha, \beta}(B(\theta, |\theta - \varphi|))}, \quad |\theta - \varphi| > 2|\theta - \theta'|,$$

$$(17) \quad \|K(\theta, \varphi) - K(\theta, \varphi')\|_{\mathbb{B}} \lesssim \frac{|\varphi - \varphi'|}{|\theta - \varphi|} \frac{1}{\mu_{\alpha, \beta}(B(\theta, |\theta - \varphi|))}, \quad |\theta - \varphi| > 2|\varphi - \varphi'|.$$

Notice that in these formulas, the ball (interval) $B(\theta, |\theta - \varphi|)$ can be replaced by $B(\varphi, |\varphi - \theta|)$, in view of the doubling property of $\mu_{\alpha, \beta}$.

We will show that the following kernels, with values in properly chosen Banach spaces \mathbb{B} , satisfy the standard estimates.

- (I) The kernel associated with the Jacobi-Poisson semigroup maximal operator,

$$\mathfrak{H}^{\alpha, \beta}(\theta, \varphi) = \{H_t^{\alpha, \beta}(\theta, \varphi)\}_{t>0}, \quad \mathbb{B} = \mathbb{X} \subset L^\infty(dt),$$

where \mathbb{X} is the closed separable subspace of $L^\infty(dt)$ consisting of all continuous functions f on $(0, \infty)$ which have finite limits as $t \rightarrow 0^+$ and as $t \rightarrow \infty$. Observe that $\{H_t^{\alpha, \beta}(\theta, \varphi)\}_{t>0} \in \mathbb{X}$, for $\theta \neq \varphi$, as can be seen from Proposition 2.3 and the bound $\mathfrak{q} \gtrsim (\theta - \varphi)^2$, and the series representation (see the proof of Lemma 3.8).

- (II) The kernels associated with Riesz-Jacobi transforms,

$$R_N^{\alpha, \beta}(\theta, \varphi) = \frac{1}{\Gamma(N)} \int_0^\infty \partial_\theta^N H_t^{\alpha, \beta}(\theta, \varphi) t^{N-1} dt, \quad \mathbb{B} = \mathbb{C},$$

where $N = 1, 2, \dots$

- (III) The kernels associated with mixed square functions,

$$\mathfrak{G}_{M, N}^{\alpha, \beta}(\theta, \varphi) = \{\partial_\theta^N \partial_t^M H_t^{\alpha, \beta}(\theta, \varphi)\}_{t>0}, \quad \mathbb{B} = L^2(t^{2M+2N-1}dt),$$

where $M, N = 0, 1, 2, \dots$ are such that $M + N > 0$.

- (IVa) The kernels associated with Laplace transform type multipliers,

$$K_\phi^{\alpha, \beta}(\theta, \varphi) = - \int_0^\infty \phi(t) \partial_t H_t^{\alpha, \beta}(\theta, \varphi) dt, \quad \mathbb{B} = \mathbb{C},$$

where $\phi \in L^\infty(dt)$.

- (IVb) The kernels associated with Laplace-Stieltjes transform type multipliers,

$$K_\nu^{\alpha, \beta}(\theta, \varphi) = \int_{(0, \infty)} H_t^{\alpha, \beta}(\theta, \varphi) d\nu(t), \quad \mathbb{B} = \mathbb{C},$$

where ν is a signed or complex Borel measure on $(0, \infty)$ with total variation $|\nu|$ satisfying

$$(18) \quad \int_{(0, \infty)} e^{-t|\frac{\alpha+\beta+1}{2}|} d|\nu|(t) < \infty.$$

When $K(\theta, \varphi)$ is scalar-valued, i.e. $\mathbb{B} = \mathbb{C}$, it is well known that the bounds (16) and (17) follow from the more convenient gradient estimate

$$(19) \quad \|\partial_\theta K(\theta, \varphi)\|_{\mathbb{B}} + \|\partial_\varphi K(\theta, \varphi)\|_{\mathbb{B}} \lesssim \frac{1}{|\theta - \varphi| \mu_{\alpha, \beta}(B(\theta, |\theta - \varphi|))}.$$

We shall see that the same holds also in the vector-valued cases we consider. Then the derivatives in (19) are taken in the weak sense, which means that for any $\mathbf{v} \in \mathbb{B}^*$

$$(20) \quad \langle \mathbf{v}, \partial_\theta K(\theta, \varphi) \rangle = \partial_\theta \langle \mathbf{v}, K(\theta, \varphi) \rangle$$

and similarly for ∂_φ . If these weak derivatives $\partial_\theta K(\theta, \varphi)$ and $\partial_\varphi K(\theta, \varphi)$ exist as elements of \mathbb{B} and their norms satisfy (19), the scalar-valued case applies and (16) and (17) follow.

The result below extends to all $\alpha, \beta > -1$ the estimates obtained in [7, Section 4] for the restricted range $\alpha, \beta \geq -1/2$. Moreover, here we also consider multipliers of Laplace and Laplace-Stieltjes transform type, which were merely mentioned in [7] and which cover as a special case the imaginary powers of $\mathcal{J}^{\alpha, \beta}$ (or $\mathcal{J}^{\alpha, \beta} \Pi_0$ when $\alpha + \beta + 1 = 0$) investigated there.

Theorem 4.1. *Let $\alpha, \beta > -1$. Then the kernels (I)-(III), (IVa) and (IVb) satisfy the standard estimates (15), (16) and (17) with \mathbb{B} as indicated above.*

In the proof we tacitly assume that passing with the differentiation in θ or φ under integrals against dt or $d\nu(t)$ is legitimate. In fact, such manipulations can easily be verified by means of the dominated convergence theorem and the estimates obtained in Corollary 3.5 and Lemma 3.8.

Proof of Theorem 4.1. We treat each of the kernels separately.

The case of $\mathfrak{H}^{\alpha, \beta}(\theta, \varphi)$. We first deal with the growth condition. Clearly, it suffices to prove independently the two bounds emerging from (15) by choosing $\mathbb{B} = L^\infty((1, \infty), dt)$ and $\mathbb{B} = L^\infty((0, 1), dt)$. These, however, are immediate consequences of Corollary 3.9 (with $M = N = L = 0, p = \infty$) and Corollary 3.5 (taken with $M = N = L = 0$) combined with Lemma 3.7 (specified to $p = \infty, s = 0$), respectively.

To obtain the smoothness estimates, we must verify that the weak derivatives $\partial_\theta \mathfrak{H}^{\alpha, \beta}(\theta, \varphi)$ and $\partial_\varphi \mathfrak{H}^{\alpha, \beta}(\theta, \varphi)$ exist in the sense of (20) and satisfy (19). In this case, \mathbf{v} is a complex measure in $[0, \infty]$, written $d\mathbf{v}$, and

$$\langle \mathbf{v}, \mathfrak{H}^{\alpha, \beta}(\theta, \varphi) \rangle = \int_{[0, \infty]} H_t^{\alpha, \beta}(\theta, \varphi) d\mathbf{v}(t).$$

It is enough to consider the derivative with respect to θ . We claim that

$$\partial_\theta \langle \mathbf{v}, \mathfrak{H}^{\alpha, \beta}(\theta, \varphi) \rangle = \int_{[0, \infty]} \partial_\theta H_t^{\alpha, \beta}(\theta, \varphi) d\mathbf{v}(t), \quad \theta \neq \varphi;$$

observe that $\{\partial_\theta H_t^{\alpha, \beta}(\theta, \varphi)\}_{t>0} \in \mathbb{X}$, $\theta \neq \varphi$, as can be seen from Proposition 2.3 and the bound $\mathfrak{q} \gtrsim (\theta - \varphi)^2$. This claim would imply that for $\theta \neq \varphi$ the weak derivative $\partial_\theta \mathfrak{H}^{\alpha, \beta}(\theta, \varphi)$ exists and equals $\{\partial_\theta H_t^{\alpha, \beta}(\theta, \varphi)\}_{t>0}$. To see that it then also satisfies (19), we first consider large t and observe that the estimate

$$\|\partial_\theta H_t^{\alpha, \beta}(\theta, \varphi)\|_{L^\infty((1, \infty), dt)} \lesssim \frac{1}{|\theta - \varphi| \mu_{\alpha, \beta}(B(\theta, |\theta - \varphi|))}, \quad \theta \neq \varphi,$$

follows from Corollary 3.9 (specified to $M = L = 0$, $N = W = 1$, $p = \infty$). For small t , we have

$$\|\partial_\theta H_t^{\alpha,\beta}(\theta, \varphi)\|_{L^\infty((0,1),dt)} \lesssim \frac{1}{|\theta - \varphi| \mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}, \quad \theta \neq \varphi,$$

in view of Corollary 3.5 (with $M = L = 0$, $N = 1$) and Lemma 3.7 (taken with $W = 1$, $p = \infty$, $s = 1$).

To prove the claim, we verify that

$$(21) \quad \int_{\theta_1}^{\theta_2} \int_{[0,\infty]} \partial_\theta H_t^{\alpha,\beta}(\theta, \varphi) dv(t) d\theta = \int_{[0,\infty]} H_t^{\alpha,\beta}(\theta_2, \varphi) dv(t) - \int_{[0,\infty]} H_t^{\alpha,\beta}(\theta_1, \varphi) dv(t),$$

where θ_1 and θ_2 are chosen in $[0, \pi]$ so that $\varphi \notin [\theta_1, \theta_2]$. Then the claim follows by differentiation with respect to θ_2 . Finally, (21) follows from Fubini's theorem, since $\partial_\theta H_t^{\alpha,\beta}(\theta, \varphi)$ is continuous and bounded in $[0, \infty] \times [\theta_1, \theta_2]$.

The case of $R_N^{\alpha,\beta}(\theta, \varphi)$. To prove the growth condition, it is enough to verify that

$$\|\partial_\theta^N H_t^{\alpha,\beta}(\theta, \varphi)\|_{L^1(t^{N-1}dt)} \lesssim \frac{1}{\mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}, \quad \theta \neq \varphi.$$

This, however, is a consequence of Corollary 3.9 (taken with $M = L = 0$, $W = N$, $p = 1$) and Corollary 3.5 (with $M = L = 0$) combined with Lemma 3.7 (specified to $W = N$, $p = 1$, $s = 0$).

In order to show the gradient bound (19), it suffices to check that

$$\left\| |\nabla_{\theta,\varphi} \partial_\theta^N H_t^{\alpha,\beta}(\theta, \varphi)| \right\|_{L^1(t^{N-1}dt)} \lesssim \frac{1}{|\theta - \varphi| \mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}, \quad \theta \neq \varphi.$$

This estimate follows by means of Corollary 3.9 (applied with $M = 0$, $p = 1$) and Corollary 3.5 (with $M = 0$) together with Lemma 3.7 (specified to $W = N$, $p = 1$, $s = 1$).

The case of $\mathfrak{S}_{M,N}^{\alpha,\beta}(\theta, \varphi)$. The growth condition is a straightforward consequence of Corollary 3.9 (with $L = 0$, $W = 2M + 2N$, $p = 2$), Corollary 3.5 (with $L = 0$) and Lemma 3.7 (taken with $W = 2M + 2N$, $p = 2$, $s = 0$).

Next, we show the gradient estimate (19), which amounts to

$$\left\| |\nabla_{\theta,\varphi} \partial_\theta^N \partial_t^M H_t^{\alpha,\beta}(\theta, \varphi)| \right\|_{L^2(t^{2M+2N-1}dt)} \lesssim \frac{1}{|\theta - \varphi| \mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}, \quad \theta \neq \varphi,$$

where $\nabla_{\theta,\varphi}$ is taken in the weak sense. This follows with the aid of Corollary 3.9 (with $W = 2M + 2N$, $p = 2$), Corollary 3.5 and Lemma 3.7 (applied with $W = 2M + 2N$, $p = 2$, $s = 1$); cf. the arguments given for the case $\mathfrak{H}^{\alpha,\beta}(\theta, \varphi)$ above.

The case of $K_\phi^{\alpha,\beta}(\theta, \varphi)$. The growth bound is a direct consequence of the assumption $\phi \in L^\infty(dt)$, Corollary 3.9 (specified to $M = 1$, $N = L = 0$, $W = 1$, $p = 1$), Corollary 3.5 (with $M = 1$, $N = L = 0$) and Lemma 3.7 (taken with $W = 1$, $p = 1$, $s = 0$).

Since ϕ is bounded, to prove the gradient estimate it is enough to verify that

$$\left\| |\nabla_{\theta,\varphi} \partial_t H_t^{\alpha,\beta}(\theta, \varphi)| \right\|_{L^1(dt)} \lesssim \frac{1}{|\theta - \varphi| \mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}, \quad \theta \neq \varphi.$$

Now applying Corollary 3.9 (with $M = 1$, $W = 1$, $p = 1$ and either $N = 1$, $L = 0$ or $N = 0$, $L = 1$), Corollary 3.5 (specified to $M = 1$ and either $N = 1$, $L = 0$ or $N = 0$, $L = 1$) and Lemma 3.7 (taken with $W = 1$, $p = 1$, $s = 1$) we arrive at the desired bound.

The case of $K_{\nu}^{\alpha,\beta}(\theta, \varphi)$. To show the growth condition it is enough, by the assumption (18) concerning the measure ν , to check that

$$\begin{aligned} \left\| e^{t|\frac{\alpha+\beta+1}{2}|} H_t^{\alpha,\beta}(\theta, \varphi) \right\|_{L^\infty((1,\infty),dt)} &\lesssim \frac{1}{\mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}, & \theta \neq \varphi, \\ \| H_t^{\alpha,\beta}(\theta, \varphi) \|_{L^\infty((0,1),dt)} &\lesssim \frac{1}{\mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}, & \theta \neq \varphi. \end{aligned}$$

The first estimate above is an immediate consequence of Corollary 3.10 (applied with $N = L = 0$). On the other hand, the remaining bound is just part of the growth condition for $\mathfrak{H}^{\alpha,\beta}(\theta, \varphi)$, which is already justified.

Taking (18) into account, to verify the gradient estimate (19), it suffices to show that

$$\begin{aligned} \left\| e^{t|\frac{\alpha+\beta+1}{2}|} |\nabla_{\theta,\varphi} H_t^{\alpha,\beta}(\theta, \varphi)| \right\|_{L^\infty((1,\infty),dt)} &\lesssim \frac{1}{|\theta - \varphi| \mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}, & \theta \neq \varphi, \\ \left\| |\nabla_{\theta,\varphi} H_t^{\alpha,\beta}(\theta, \varphi)| \right\|_{L^\infty((0,1),dt)} &\lesssim \frac{1}{|\theta - \varphi| \mu_{\alpha,\beta}(B(\theta, |\theta - \varphi|))}, & \theta \neq \varphi. \end{aligned}$$

Again, an application of Corollary 3.10 (with either $N = 1, L = 0$ or $N = 0, L = 1$) produces the first bound. The second one is contained in the proof of the gradient estimate for $\mathfrak{H}^{\alpha,\beta}(\theta, \varphi)$.

The proof of Theorem 4.1 is complete. \square

5. CALDERÓN-ZYGMUND OPERATORS

Let \mathbb{B} be a Banach space and suppose that T is a linear operator assigning to each $f \in L^2(d\mu_{\alpha,\beta})$ a strongly measurable \mathbb{B} -valued function Tf on $[0, \pi]$. Then T is said to be a (vector-valued) Calderón-Zygmund operator in the sense of the space $([0, \pi], d\mu_{\alpha,\beta}, |\cdot|)$ associated with \mathbb{B} if

- (A) T is bounded from $L^2(d\mu_{\alpha,\beta})$ to $L_{\mathbb{B}}^2(d\mu_{\alpha,\beta})$,
- (B) there exists a standard \mathbb{B} -valued kernel $K(\theta, \varphi)$ such that

$$Tf(\theta) = \int_0^\pi K(\theta, \varphi) f(\varphi) d\mu_{\alpha,\beta}(\varphi), \quad \text{a.a. } \theta \notin \text{supp } f,$$

for $f \in L^\infty([0, \pi])$.

Here integration of \mathbb{B} -valued functions is understood in Bochner's sense, and $L_{\mathbb{B}}^2(d\mu_{\alpha,\beta})$ is the Bochner-Lebesgue space of all \mathbb{B} -valued $d\mu_{\alpha,\beta}$ -square integrable functions on $[0, \pi]$.

It is well known that a large part of the classical theory of Calderón-Zygmund operators remains valid, with appropriate adjustments, when the underlying space is of homogeneous type and the associated kernels are vector-valued, see for instance [10, 11]. In particular, if T is a Calderón-Zygmund operator in the sense of $([0, \pi], d\mu_{\alpha,\beta}, |\cdot|)$ associated with a Banach space \mathbb{B} , then its mapping properties in weighted L^p spaces follow from the general theory.

Let

$$\mathcal{H}_t^{\alpha,\beta} f(\theta) = \int_0^\pi H_t^{\alpha,\beta}(\theta, \varphi) f(\varphi) d\mu_{\alpha,\beta}(\varphi), \quad t > 0, \quad \theta \in [0, \pi],$$

be the Jacobi-Poisson semigroup. For $\alpha, \beta > -1$ consider the following operators defined initially in $L^2(d\mu_{\alpha,\beta})$.

- (I) The Jacobi-Poisson semigroup maximal operator

$$\mathcal{H}_*^{\alpha,\beta} f = \left\| \mathcal{H}_t^{\alpha,\beta} f \right\|_{L^\infty(dt)}.$$

(II) Riesz-Jacobi transforms of orders $N = 1, 2, \dots$

$$R_N^{\alpha,\beta} f = \sum_{n=1}^{\infty} \left| n + \frac{\alpha + \beta + 1}{2} \right|^{-N} \langle f, \mathcal{P}_n^{\alpha,\beta} \rangle_{d\mu_{\alpha,\beta}} \partial_{\theta}^N \mathcal{P}_n^{\alpha,\beta},$$

where $\langle f, \mathcal{P}_n^{\alpha,\beta} \rangle_{d\mu_{\alpha,\beta}}$ are the Fourier-Jacobi coefficients of f .

(III) Littlewood-Paley-Stein type mixed square functions

$$g_{M,N}^{\alpha,\beta}(f) = \left\| \partial_{\theta}^N \partial_t^M \mathcal{H}_t^{\alpha,\beta} f \right\|_{L^2(t^{2M+2N-1} dt)},$$

where $M, N = 0, 1, 2, \dots$ and $M + N > 0$.

(IV) Multipliers of Laplace and Laplace-Stieltjes transform type

$$M_{\mathbf{m}}^{\alpha,\beta} f = \sum_{n=0}^{\infty} \mathbf{m} \left(\left| n + \frac{\alpha + \beta + 1}{2} \right| \right) \langle f, \mathcal{P}_n^{\alpha,\beta} \rangle_{d\mu_{\alpha,\beta}} \mathcal{P}_n^{\alpha,\beta},$$

where either $\mathbf{m}(z) = \int_0^{\infty} z e^{-tz} \phi(t) dt$ with $\phi \in L^{\infty}(dt)$ or $\mathbf{m}(z) = \int_{(0,\infty)} e^{-tz} d\nu(t)$ with ν being a signed or complex Borel measure on $(0, \infty)$ whose total variation satisfies (18).

The formulas defining $\mathcal{H}_*^{\alpha,\beta}$ and $g_{M,N}^{\alpha,\beta}$ are understood pointwise and are actually valid for general functions f from weighted L^p spaces with Muckenhoupt weights. This is because for such f the integral defining $\mathcal{H}_t^{\alpha,\beta} f(\theta)$ is well defined and produces a smooth function of $(t, \theta) \in (0, \infty) \times [0, \pi]$, see [7, Section 2]. The series defining $R_N^{\alpha,\beta}$ and $M_{\mathbf{m}}^{\alpha,\beta}$ indeed converge in $L^2(d\mu_{\alpha,\beta})$, which is clear in the case of $M_{\mathbf{m}}^{\alpha,\beta}$, since the values of \mathbf{m} that occur here stay bounded. For $R_N^{\alpha,\beta}$ the convergence follows by [7, Lemma 3.1], see the proof of [7, Proposition 2.2] in the case of $R_N^{\alpha,\beta}$.

As a consequence of Theorem 4.1 we get the following result.

Theorem 5.1. *Assume that $\alpha, \beta > -1$. The Riesz-Jacobi transforms and the multipliers of Laplace and Laplace-Stieltjes transform type are scalar-valued Calderón-Zygmund operators in the sense of the space $([0, \pi], d\mu_{\alpha,\beta}, |\cdot|)$. Furthermore, the Jacobi-Poisson semigroup maximal operator and the mixed square functions can be viewed as vector-valued Calderón-Zygmund operators in the sense of $([0, \pi], d\mu_{\alpha,\beta}, |\cdot|)$, associated with the Banach spaces $\mathbb{B} = \mathbb{X}$ and $\mathbb{B} = L^2(t^{2M+2N-1} dt)$, respectively.*

Proof. The standard estimates are provided in all the cases by Theorem 4.1. Thus it suffices to verify L^2 boundedness and kernel associations (conditions (A) and (B) above). This, however, was essentially done in [7, Section 3], since the arguments given there are actually valid for all $\alpha, \beta > -1$ if combined with the estimates proved (in some cases implicitly) in Section 4. An exception here are the Laplace and Laplace-Stieltjes type multipliers. But in these cases the boundedness in L^2 is straightforward, and kernel associations are justified according to the outline opening the proof of [7, Proposition 2.3], see [7, Section 3, pp. 732–733]. Since all the necessary ingredients are contained in [7] and in the present paper, we leave further details to interested readers. \square

Denote by $A_p^{\alpha,\beta}$, $1 \leq p < \infty$, the Muckenhoupt classes of weights related to the space $([0, \pi], d\mu_{\alpha,\beta}, |\cdot|)$ (see [7, Section 1] for the definition).

Corollary 5.2. *Let $\alpha, \beta > -1$. The Riesz-Jacobi transforms and the multipliers of Laplace and Laplace-Stieltjes type extend to bounded linear operators on $L^p(w d\mu_{\alpha,\beta})$, $w \in A_p^{\alpha,\beta}$, $1 < p < \infty$,*

and from $L^1(wd\mu_{\alpha,\beta})$ to weak $L^1(wd\mu_{\alpha,\beta})$, $w \in A_1^{\alpha,\beta}$. The same boundedness properties hold for the Jacobi-Poisson semigroup maximal operator and the mixed square functions, viewed as scalar-valued sublinear operators.

Proof. The part concerning $R_N^{\alpha,\beta}$ and $M_m^{\alpha,\beta}$ is a direct consequence of Theorem 5.1 and the general theory. The remaining part follows by Theorem 5.1 and the arguments given in the proof of [7, Corollary 2.5]. \square

6. EXACT BEHAVIOR OF THE JACOBI-POISSON KERNEL

We give another application of the representations in Proposition 2.3, which is interesting and important in its own right. We will describe in a sharp way the behavior of the kernels $\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi)$ and $H_t^{\alpha,\beta}(\theta, \varphi)$. The result below extends sharp estimates for the Jacobi-Poisson kernel obtained in [8, Theorem 5.2] under the restriction $\alpha, \beta \geq -1/2$.

Theorem 6.1. *Let $\alpha, \beta > -1$. Then*

$$\begin{aligned} H_t^{\alpha,\beta}(\theta, \varphi) &\simeq \mathbb{H}_t^{\alpha,\beta}(\theta, \varphi) \\ &\simeq \left(t^2 + \theta^2 + \varphi^2\right)^{-\alpha-1/2} \left(t^2 + (\pi - \theta)^2 + (\pi - \varphi)^2\right)^{-\beta-1/2} \frac{t}{t^2 + (\theta - \varphi)^2}, \end{aligned}$$

uniformly in $0 < t \leq 1$ and $\theta, \varphi \in [0, \pi]$, and

$$H_t^{\alpha,\beta}(\theta, \varphi) \simeq \exp\left(-t \frac{|\alpha + \beta + 1|}{2}\right), \quad \mathbb{H}_t^{\alpha,\beta}(\theta, \varphi) \simeq \exp\left(-t \frac{\alpha + \beta + 1}{2}\right),$$

uniformly in $t \geq 1$ and $\theta, \varphi \in [0, \pi]$.

To prove this we will need some technical results, one of which is Lemma 3.2 (a). Note that this lemma remains true if the integration is restricted to the subinterval $(1/2, 1]$. This follows from the structure of $d\Pi_\nu$ and the fact that the integrand is positive and increasing.

Lemma 6.2. *Let $\tau > 0$ be fixed. Then*

$$\frac{1}{a^\tau} - \frac{1}{b^\tau} - \frac{1}{c^\tau} + \frac{1}{d^\tau} \gtrsim \frac{(b \wedge c - a)^2 \wedge a^2}{a^{\tau+2}},$$

uniformly in $0 < a \leq b, c \leq d$ satisfying $a + d = b + c$.

Proof. We can assume that $b \leq c$. Then the right-hand side is independent of c and d . In the left-hand side, we therefore replace c and d by $c + s$ and $d + s$, respectively, where $s \geq b - c$. By differentiating, we see that the function $s \mapsto -(c + s)^{-\tau} + (d + s)^{-\tau}$ is increasing. As a result, we need only consider the extreme case $s = b - c$, which means proving the lemma for $b = c$.

Writing $h = b - a$, and letting $f(x) = x^{-\tau}$, the left-hand side is now the second difference $f(a) - 2f(a + h) + f(a + 2h)$, which equals $f''(\xi)h^2$ for some $\xi \in (a, a + 2h)$. Now if $h > Ca$ for some large $C = C(\tau)$, the inequality of the lemma is trivial, since the term $a^{-\tau}$ will dominate in the left-hand side. But if $h \leq Ca$, we have $f''(\xi) \simeq a^{-\tau-2}$, and the conclusion follows again. \square

Let $\sigma > 1$ be fixed. Then one easily verifies that

$$(22) \quad |x^{-\sigma} - y^{-\sigma}| \simeq \frac{|x - y|}{(x \vee y)(x \wedge y)^\sigma}, \quad x, y > 0.$$

Proof of Theorem 6.1. We first prove the estimates for $\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi)$. Among the four ranges of the type parameters distinguished in Proposition 2.3, it is enough to consider only two. Indeed, when $\alpha, \beta \geq -1/2$ the desired bounds are contained in [8, Theorem 5.2], and the cases $\beta < -1/2 \leq \alpha$ and $\alpha < -1/2 \leq \beta$ are essentially the same. In what follows we denote for $t > 0$ and $\theta, \varphi \in [0, \pi]$

$$X := \frac{\sin \frac{\theta}{2} \sin \frac{\varphi}{2}}{\cosh \frac{t}{2} - \cos \frac{\theta}{2} \cos \frac{\varphi}{2}}, \quad Y := \frac{\cos \frac{\theta}{2} \cos \frac{\varphi}{2}}{\cosh \frac{t}{2} - \sin \frac{\theta}{2} \sin \frac{\varphi}{2}},$$

and

$$Z := \frac{\sinh \frac{t}{2}}{\left(\cosh \frac{t}{2} - \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right)^{\alpha+1/2} \left(\cosh \frac{t}{2} - \sin \frac{\theta}{2} \sin \frac{\varphi}{2} \right)^{\beta+1/2} \left(\cosh \frac{t}{2} - \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right)}.$$

Notice that $0 \leq X, Y < 1$ and that Z is comparable, uniformly in $0 < t \leq 1$ and $\theta, \varphi \in [0, \pi]$, with the expression describing the short-time behavior in Theorem 6.1; see the proof of [8, Theorem 5.2]. Moreover, Z has the same long-time behavior as that asserted for $\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi)$. Thus that part of the statement of Theorem 6.1 which deals with $\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi)$ can be written simply as

$$(23) \quad \mathbb{H}_t^{\alpha,\beta}(\theta, \varphi) \simeq Z, \quad t > 0, \quad \theta, \varphi \in [0, \pi].$$

Case 1: $-1 < \alpha < -1/2 \leq \beta$. By Proposition 2.3,

$$\begin{aligned} \mathbb{H}_t^{\alpha,\beta}(\theta, \varphi) &= \iint -\partial_u \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) \Pi_\alpha(u) du d\Pi_\beta(v) \\ &\quad + \iint \Psi^{\alpha,\beta}(t, \theta, \varphi, u, v) d\Pi_{-1/2}(u) d\Pi_\beta(v) \\ &\equiv I_1 + I_2. \end{aligned}$$

One finds that the integral I_1 is dominated (up to a multiplicative constant) by its restriction to the subsquare $(1/2, 1]^2$ and that the essential contribution to I_2 comes from integrating over $(1/2, 1]^2$. In view of Lemma 2.2, the measures $|\Pi_\alpha(u)| du$ and $d\Pi_{\alpha+1}$ are comparable on $(1/2, 1]$, and we infer that

$$\begin{aligned} I_1 &\lesssim \sinh \frac{t}{2} \sin \frac{\theta}{2} \sin \frac{\varphi}{2} \iint \frac{d\Pi_{\alpha+1}(u) d\Pi_\beta(v)}{\left(\cosh \frac{t}{2} - u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right)^{\alpha+\beta+3}}, \\ I_2 &\simeq \sinh \frac{t}{2} \int \frac{d\Pi_\beta(v)}{\left(\cosh \frac{t}{2} - \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right)^{\alpha+\beta+2}}, \end{aligned}$$

uniformly in $t > 0$ and $\theta, \varphi \in [0, \pi]$. Applying now Lemma 3.2 (a) to I_1 twice, first to the integral against $d\Pi_\beta$, with the parameters $\nu = \beta$, $\kappa = 0$, $\gamma = \alpha + \beta + 3$, $A = \cosh \frac{t}{2} - u \sin \frac{\theta}{2} \sin \frac{\varphi}{2}$, $B = \cos \frac{\theta}{2} \cos \frac{\varphi}{2}$, and then to the resulting integral against $d\Pi_{\alpha+1}$, with the parameters $\nu = \alpha + 1$, $\kappa = \beta + 1/2$, $\gamma = \alpha + 5/2$, $D = \cosh \frac{t}{2}$, $A = \cosh \frac{t}{2} - \cos \frac{\theta}{2} \cos \frac{\varphi}{2}$, $B = \sin \frac{\theta}{2} \sin \frac{\varphi}{2}$, we arrive at the bound

$$I_1 \lesssim XZ.$$

Applying once again Lemma 3.2 (a), this time to I_2 and with the parameters $\nu = \beta$, $\kappa = 0$, $\gamma = \alpha + \beta + 2$, $A = \cosh \frac{t}{2} - \sin \frac{\theta}{2} \sin \frac{\varphi}{2}$, $B = \cos \frac{\theta}{2} \cos \frac{\varphi}{2}$, we get

$$I_2 \simeq (1 - X)^{-\alpha-1/2} Z.$$

Estimating I_1 from below is slightly more subtle. Notice that

$$I_1 = \sum_{\eta=\pm 1} \iint_{(0,1]^2} (\partial_u \Psi^{\alpha,\beta}(t, \theta, \varphi, u, \eta v) - \partial_u \Psi^{\alpha,\beta}(t, \theta, \varphi, -u, \eta v)) |\Pi_\alpha(u)| du d\Pi_\beta(v);$$

here the integrand in each double integral is nonnegative, and the one corresponding to $\eta = 1$ is dominating. Thus restricting the set of integration to $(1/2, 1]^2$ and making use of Lemma 2.2, we write

$$\begin{aligned} I_1 &\gtrsim \iint_{(1/2, 1]^2} (\partial_u \Psi^{\alpha, \beta}(t, \theta, \varphi, u, v) - \partial_u \Psi^{\alpha, \beta}(t, \theta, \varphi, -u, v)) d\Pi_{\alpha+1}(u) d\Pi_{\beta}(v) \\ &\simeq \sinh \frac{t}{2} \sin \frac{\theta}{2} \sin \frac{\varphi}{2} \iint_{(1/2, 1]^2} \left[\frac{1}{\left(\cosh \frac{t}{2} - u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right)^{\alpha+\beta+3}} \right. \\ &\quad \left. - \frac{1}{\left(\cosh \frac{t}{2} + u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right)^{\alpha+\beta+3}} \right] d\Pi_{\alpha+1}(u) d\Pi_{\beta}(v). \end{aligned}$$

Applying (22) to the expression in square brackets above, we get

$$\begin{aligned} I_1 &\gtrsim \iint_{(1/2, 1]^2} \frac{\sinh \frac{t}{2} (\sin \frac{\theta}{2} \sin \frac{\varphi}{2})^2 u d\Pi_{\alpha+1}(u) d\Pi_{\beta}(v)}{\left(\cosh \frac{t}{2} + u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right) \left(\cosh \frac{t}{2} - u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right)^{\alpha+\beta+3}} \\ &\gtrsim \iint_{(1/2, 1]^2} \frac{\sinh \frac{t}{2} (\sin \frac{\theta}{2} \sin \frac{\varphi}{2})^2 d\Pi_{\alpha+1}(u) d\Pi_{\beta}(v)}{\left(\cosh \frac{t}{2} + \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right) \left(\cosh \frac{t}{2} - u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right)^{\alpha+\beta+3}}. \end{aligned}$$

The last integral is comparable with an analogous integral over the larger square $[-1, 1]^2$, see the comment following Theorem 6.1. Using now Lemma 3.2 (a) twice, first for the integral against $d\Pi_{\beta}$ (with the parameters $\nu = \beta$, $\kappa = 1$, $\gamma = \alpha + \beta + 3$, $D = \cosh \frac{t}{2} + \sin \frac{\theta}{2} \sin \frac{\varphi}{2}$, $A = \cosh \frac{t}{2} - u \sin \frac{\theta}{2} \sin \frac{\varphi}{2}$, $B = \cos \frac{\theta}{2} \cos \frac{\varphi}{2}$) and then with the resulting integral against $d\Pi_{\alpha+1}$ (with $\nu = \alpha + 1$, $\kappa = \beta + 1/2$, $\gamma = \alpha + 5/2$, $D = \cosh \frac{t}{2}$, $A = \cosh \frac{t}{2} - \cos \frac{\theta}{2} \cos \frac{\varphi}{2}$, $B = \sin \frac{\theta}{2} \sin \frac{\varphi}{2}$), we arrive at the bound

$$I_1 \gtrsim \frac{X^2}{X+1} Z \simeq X^2 Z.$$

Summing up, we have proved that

$$X^2 Z + (1 - X)^{-\alpha-1/2} Z \lesssim \mathbb{H}_t^{\alpha, \beta}(\theta, \varphi) \lesssim X Z + (1 - X)^{-\alpha-1/2} Z,$$

uniformly in $t > 0$ and $\theta, \varphi \in [0, \pi]$, and (23) follows.

Case 2: $-1 < \alpha, \beta < -1/2$. In view of Proposition 2.3,

$$\begin{aligned} \mathbb{H}_t^{\alpha, \beta}(\theta, \varphi) &= \iint \partial_u \partial_v \Psi^{\alpha, \beta}(t, \theta, \varphi, u, v) \Pi_{\alpha}(u) du \Pi_{\beta}(v) dv \\ &\quad + \iint -\partial_u \Psi^{\alpha, \beta}(t, \theta, \varphi, u, v) \Pi_{\alpha}(u) du d\Pi_{-1/2}(v) \\ &\quad + \iint -\partial_v \Psi^{\alpha, \beta}(t, \theta, \varphi, u, v) d\Pi_{-1/2}(u) \Pi_{\beta}(v) dv \\ &\quad + \iint \Psi^{\alpha, \beta}(t, \theta, \varphi, u, v) d\Pi_{-1/2}(u) d\Pi_{-1/2}(v) \\ &\equiv J_1 + J_2 + J_3 + J_4. \end{aligned}$$

Clearly, the main contribution to J_4 comes from the point $(u, v) = (1, 1)$, and so

$$J_4 \simeq \Psi^{\alpha, \beta}(t, \theta, \varphi, 1, 1) \simeq (1 - X)^{-\alpha-1/2} (1 - Y)^{-\beta-1/2} Z \leq Z.$$

To bound the remaining integrals from above, we proceed as in Case 1, obtaining

$$J_1 \lesssim \iint \partial_u \partial_v \Psi^{\alpha, \beta}(t, \theta, \varphi, u, v) d\Pi_{\alpha+1}(u) d\Pi_{\beta+1}(v),$$

$$J_2 \lesssim \int \partial_u \Psi^{\alpha, \beta}(t, \theta, \varphi, u, 1) d\Pi_{\alpha+1}(u),$$

$$J_3 \lesssim \int \partial_v \Psi^{\alpha, \beta}(t, \theta, \varphi, 1, v) d\Pi_{\beta+1}(v).$$

Then applying repeatedly Lemma 3.2 (a) with suitably chosen parameters, we get

$$J_1 \lesssim XYZ \leq Z, \quad J_2 \lesssim X(1-Y)^{-\beta-1/2}Z \leq Z, \quad J_3 \lesssim (1-X)^{-\alpha-1/2}YZ \leq Z.$$

To estimate J_2 and J_3 from below, we use the same trick as for I_1 in Case 1. By means of Lemma 2.2 and (22), we can write

$$J_2 \gtrsim \frac{\sinh \frac{t}{2} \left(\sin \frac{\theta}{2} \sin \frac{\varphi}{2} \right)^2}{\cosh \frac{t}{2} + \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - \cos \frac{\theta}{2} \cos \frac{\varphi}{2}} \int_{(1/2, 1]} \frac{d\Pi_{\alpha+1}(u)}{\left(\cosh \frac{t}{2} - u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right)^{\alpha+\beta+3}}.$$

Then Lemma 3.2 (a) shows that

$$J_2 \gtrsim \frac{X^2}{X+1} (1-Y)^{-\beta-1/2} Z \simeq X^2 (1-Y)^{-\beta-1/2} Z.$$

The case of J_3 is parallel, we have

$$J_3 \gtrsim (1-X)^{-\alpha-1/2} \frac{Y^2}{Y+1} Z \simeq (1-X)^{-\alpha-1/2} Y^2 Z.$$

Finally, we focus on the more delicate integral J_1 . Observe that

$$J_1 = \iint \sum_{\xi, \eta = \pm 1} \xi \eta \partial_u \partial_v \Psi^{\alpha, \beta}(t, \theta, \varphi, \xi u, \eta v) |\Pi_{\alpha}(u)| |\Pi_{\beta}(v)| du dv.$$

Restricting here the region of integration (the integrand is nonnegative, as we shall see in a moment) and using Lemma 2.2, we conclude

$$J_1 \gtrsim \sinh \frac{t}{2} \sin \frac{\theta}{2} \sin \frac{\varphi}{2} \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \iint_{(1/2, 1]^2} \left(\frac{1}{a^{\tau}} - \frac{1}{b^{\tau}} - \frac{1}{c^{\tau}} + \frac{1}{d^{\tau}} \right) d\Pi_{\alpha+1}(u) d\Pi_{\beta+1}(v),$$

where $\tau = \alpha + \beta + 4$, $a = \cosh \frac{t}{2} - u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2}$, $b = \cosh \frac{t}{2} - u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} + v \cos \frac{\theta}{2} \cos \frac{\varphi}{2}$, $c = \cosh \frac{t}{2} + u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2}$, $d = \cosh \frac{t}{2} + u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} + v \cos \frac{\theta}{2} \cos \frac{\varphi}{2}$.

Applying now Lemma 6.2, we get

$$J_1 \gtrsim \sinh \frac{t}{2} \sin \frac{\theta}{2} \sin \frac{\varphi}{2} \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \iint_{(1/2, 1]^2} \frac{(b \wedge c - a)^2 \wedge a^2}{a^{\alpha+\beta+6}} d\Pi_{\alpha+1}(u) d\Pi_{\beta+1}(v).$$

Since

$$b \wedge c - a = 2u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} \wedge 2v \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \geq \sin \frac{\theta}{2} \sin \frac{\varphi}{2} \wedge \cos \frac{\theta}{2} \cos \frac{\varphi}{2}, \quad u, v \geq 1/2,$$

we can write

$$J_1 \gtrsim \sinh \frac{t}{2} \sin \frac{\theta}{2} \sin \frac{\varphi}{2} \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \iint_{(1/2, 1]^2} \frac{d\Pi_{\alpha+1}(u) d\Pi_{\beta+1}(v)}{\left(\cosh \frac{t}{2} - u \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - v \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right)^{\alpha+\beta+6}} \\ \times \left[\left(\cosh \frac{t}{2} - \sin \frac{\theta}{2} \sin \frac{\varphi}{2} - \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right) \wedge \sin \frac{\theta}{2} \sin \frac{\varphi}{2} \wedge \cos \frac{\theta}{2} \cos \frac{\varphi}{2} \right]^2.$$

Combining this with Lemma 3.2 (a), we see that

$$J_1 \gtrsim XY \left[1 \wedge \left(\frac{X}{1-X} \right)^2 \wedge \left(\frac{Y}{1-Y} \right)^2 \right] Z \geq (X \wedge Y)^4 Z.$$

Altogether, the above considerations justify the estimates

$$\begin{aligned} & \left((X \wedge Y)^4 + X^2(1-Y)^{-\beta-1/2} + (1-X)^{-\alpha-1/2}Y^2 + (1-X)^{-\alpha-1/2}(1-Y)^{-\beta-1/2} \right) Z \\ & \lesssim \mathbb{H}_t^{\alpha,\beta}(\theta, \varphi) \lesssim Z, \end{aligned}$$

which hold uniformly in $t > 0$ and $\theta, \varphi \in [0, \pi]$. From this, (23) follows.

We pass to the Jacobi-Poisson kernel $H_t^{\alpha,\beta}(\theta, \varphi)$. Here we can assume that $\lambda := \alpha + \beta + 1 < 0$, since otherwise the kernels $\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi)$ and $H_t^{\alpha,\beta}(\theta, \varphi)$ coincide. Then

$$H_t^{\alpha,\beta}(\theta, \varphi) = \mathbb{H}_t^{\alpha,\beta}(\theta, \varphi) + 2^{\lambda+1} c_{\alpha,\beta} \sinh \frac{\lambda t}{2}.$$

The second term here is negative for $t > 0$, so $H_t^{\alpha,\beta}(\theta, \varphi) < \mathbb{H}_t^{\alpha,\beta}(\theta, \varphi)$. Taking (23) into account, we obtain the short-time upper bound for $H_t^{\alpha,\beta}(\theta, \varphi)$. Thus what remains to show is the lower bound and the long-time upper bound for $H_t^{\alpha,\beta}(\theta, \varphi)$.

We first claim that the lower short-time bound holds provided that $t > 0$ is small enough. In view of the already justified estimates for $\mathbb{H}_t^{\alpha,\beta}(\theta, \varphi)$, this will follow once we check that

$$-2^{\lambda+1} c_{\alpha,\beta} \sinh \frac{\lambda t}{2} \leq c \mathbb{H}_t^{\alpha,\beta}(\theta, \varphi), \quad 0 < t \leq T_0,$$

for some $T_0 > 0$ and some $c < 1$. Notice that the hypergeometric series defining F_4 in (2) has nonnegative terms and that the zero-order term is 1. Thus for $t > 0$ and $\theta, \varphi \in [0, \pi]$

$$\left(\frac{2}{e} \right)^{\lambda+1} \mathbb{H}_t^{\alpha,\beta}(\theta, \varphi) + 2^{\lambda+1} c_{\alpha,\beta} \sinh \frac{\lambda t}{2} \geq \left(\frac{2}{e} \right)^{\lambda+1} c_{\alpha,\beta} \left(\frac{\sinh \frac{t}{2}}{(\cosh \frac{t}{2})^{\lambda+1}} + e^{\lambda+1} \sinh \frac{\lambda t}{2} \right).$$

Now it suffices to ensure that, given $\lambda \in (-1, 0)$, the function

$$h(s) = \frac{\sinh s}{(\cosh s)^{\lambda+1}} + e^{\lambda+1} \sinh(\lambda s)$$

satisfies $h(0) = 0$ and $h'(0) > 0$. This, however, is straightforward. The claim follows.

Next we show that the upper long-time bound for $H_t^{\alpha,\beta}(\theta, \varphi)$ holds for $t \geq 1$ and that the lower counterpart is also true provided that $t \geq T_1$ with T_1 chosen large enough. From the series representation,

$$H_t^{\alpha,\beta}(\theta, \varphi) = 2^\lambda c_{\alpha,\beta} e^{-t|\lambda|/2} + \sum_{n=1}^{\infty} e^{-t(n+\lambda/2)} \mathcal{P}_n^{\alpha,\beta}(\theta) \mathcal{P}_n^{\alpha,\beta}(\varphi).$$

The last series can be controlled by means of the bound $|\mathcal{P}_n^{\alpha,\beta}(\theta)| \lesssim n$, $n \geq 1$, see (14). More precisely, we have

$$\left| \sum_{n=1}^{\infty} e^{-t(n+\lambda/2)} \mathcal{P}_n^{\alpha,\beta}(\theta) \mathcal{P}_n^{\alpha,\beta}(\varphi) \right| \lesssim e^{-t/2} \sum_{n=1}^{\infty} n^2 e^{-t(n+\frac{\alpha+\beta}{2})} \lesssim e^{-t/2}, \quad t \geq 1.$$

Since $\alpha + \beta > -2$ and $|\lambda| < 1$, the conclusion follows.

To deal finally with the lower bound in the range $T_0 \leq t \leq T_1$, we use the semigroup property of $H_t^{\alpha,\beta}$. For $T_0 \leq t \leq 2T_0$, we have

$$H_t^{\alpha,\beta}(\theta, \varphi) = \int_0^\pi H_{t/2}^{\alpha,\beta}(\theta, \psi) H_{t/2}^{\alpha,\beta}(\psi, \varphi) d\mu_{\alpha,\beta}(\psi).$$

Since $H_{t/2}^{\alpha,\beta}(\theta, \varphi) \gtrsim 1$ in $[T_0, 2T_0] \times [0, \pi]^2$ by the above, we conclude that also $H_t^{\alpha,\beta}(\theta, \varphi)$ has a positive lower bound in the same set. In a finite number of similar steps, we will reach $t = T_1$.

The proof of Theorem 6.1 is complete. □

REFERENCES

- [1] G.E. Andrews, R. Askey, R. Roy, *Special functions*, Encyclopedia of Mathematics and its Applications, vol. 71, Cambridge University Press, Cambridge, 1999.
- [2] A.J. Castro, T.Z. Szarek, *Calderón-Zygmund operators in the Bessel setting for all possible type indices*, preprint 2011. [arXiv:1110.5659v1](#)
- [3] A. Erdélyi, W. Magnus, F. Oberhettinger, F.G. Tricomi, *Higher transcendental functions. Vol. I. Based on notes left by Harry Bateman*. Reprint of the 1953 original. Robert E. Krieger Publishing Co., Inc., Melbourne, Fla., 1981.
- [4] A. Dijksma, T.K. Koornwinder, *Spherical harmonics and the product of two Jacobi polynomials*, Indag. Math. 33 (1971), 171–196.
- [5] W.P. Johnson, *The curious history of Faà di Bruno’s formula*, Amer. Math. Monthly 109 (2002), 217–234.
- [6] B. Muckenhoupt, E.M. Stein, *Classical expansions and their relation to conjugate harmonic functions*, Trans. Amer. Math. Soc. 118 (1965), 17–92.
- [7] A. Nowak, P. Sjögren, *Calderón-Zygmund operators related to Jacobi expansions*, J. Fourier Anal. Appl. 18 (2012), 717–749.
- [8] A. Nowak, P. Sjögren, *Sharp estimates of the Jacobi heat kernel*, preprint 2011. [arXiv:1111.3145v1](#)
- [9] A. Nowak, T.Z. Szarek, *Calderón-Zygmund operators related to Laguerre function expansions of convolution type*, J. Math. Anal. Appl. 388 (2012), 801–816.
- [10] J.L. Rubio de Francia, F. Ruiz, J.L. Torrea, *Calderón-Zygmund theory for operator-valued kernels*, Adv. Math. 62 (1986), 7–48.
- [11] F. Ruiz, J.L. Torrea, *Vector-valued Calderón-Zygmund theory and Carleson measures on spaces of homogeneous nature*, Studia Math. 88 (1988), 221–243.
- [12] G. Szegő, *Orthogonal Polynomials*, 4th edn. Amer. Math. Soc. Colloq. Publ. vol. 23. Am. Math. Soc., Providence (1975).
- [13] B. Wróbel, *Multivariate spectral multipliers for tensor product orthogonal expansions*, Monatsh. Math. 168 (2012), 125–149.

ADAM NOWAK,

INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES,
ŚNIADECKICH 8, 00–956 WARSZAWA, POLAND

E-mail address: anowak@impan.pl

PETER SJÖGREN,

MATHEMATICAL SCIENCES, UNIVERSITY OF GOTHENBURG
MATHEMATICAL SCIENCES, CHALMERS UNIVERSITY OF TECHNOLOGY
SE-412 96 GÖTEBORG, SWEDEN

E-mail address: peters@chalmers.se

TOMASZ Z. SZAREK,

INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES,
ŚNIADECKICH 8, 00–956 WARSZAWA, POLAND

E-mail address: szarektomaszz@gmail.com